

WHERE TO APPLY (continued)

FLOWER GARDENS

Apply to soil after setting out plants or after flowers are well established. Miller's WEEDER is specifically recommended for weed control with the following:

Ageratum	Fernleaf Yarrow	Orpine
Alyssum	Feverfew	Proniss
Aster	Forget-me-not	Petunia
Babysbreath	Four-O'clock	Pink Heath
Bellflower	Foxglove	Poker Plant
Bleedingheart	Gaillardia	Purple Coneflower
Blood leaf	Geranium	Rose Moss
Bluefloss	Gladiolus	Salvia
Candle Larkspur	Gold Dust	Scarlet Sage
Candytufts	Golden Marguerite	Snapdragon
Chrysanthemum	Golden Tuft	Soldierwort
Cinquefoil	Iris	Strawflower
Columbine	Lavendercotton	Sunflower
Coleus	Lily	Sweet Pea
Coreopsis	Lupine	Verbena
Coral Bells	Marigold	Violet
Cosmos	Morning Glory	Wormwood
Cypripedium	Mother-of-Thyme	Zinnia
Dahlia	Mourning Bride	
Delphinium	Mourning Peaks	
Evening Primrose	Nasturtium	

ORNAMENTAL PLANTINGS

Apply to the soil in early spring. Late summer applications may prove beneficial for control of fall-germinating weeds. For newly transplanted ornamentals, apply the granules immediately after setting out whether transplanted in spring or fall. Miller's WEEDER is specifically recommended for weed control with the following:

Abelia	Fir	Pine
Andromeda, Japanese	Forsythia	Pittosporum
Arbutus	Gum	Podocarpus
Ash	Hawthorn	Poplar
Azalea	Holly	Privet
Barberry	Honeysuckle	Privet, Variegated
Boxwood	Hydrangea	Redbud
Birch	Ivy	Rhododendron
Camellia	(Baltic, Boston English & Wilson)	Rose
Chestnut	Juniper	Russian Olive
Cotoneaster	Lantana	Spiraea
Cottonwood	Lilac	Spruce
Crab Apple (Ornamental)	Locust	Sycamore
Cypress	Magnolia	Tree Peony
Deutzia	Maple	Tulip Tree
Dogwood	Mock Orange	Viburnum
Elaeagnus	Mountain Laurel	Walnut
Elm	Oak	Weigelia
Euonymus	Pachistima	Willow
	Pachysandra	Yew

DO NOT USE on the following plants:

Bugle Weed	Mesembryanthemum
Button Pink	Pansy
Carnation	Phlox
Cam	Sweet William
Germander	Telanthera

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-4H

Distributed by:

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

NET WT.
1 LB.

\$179

Weeder

*FOR WEED CONTROL
IN VEGETABLE AND
FLOWER GARDENS*

Please read entire label.
Use only as directed, noting all cautions.

Miller's WEEDER is a granular product for use as a pre-emergence application. It kills weed seeds as they germinate, before they emerge from the soil.

HOW TO APPLY

Cultivate the area to be treated so that all weeds are removed, leaving a uniformly smooth soil surface. Measure off the area to be treated... one pound of Miller's WEEDER will treat 250 sq. ft.—10' x 25' overall, or in a band 1 foot wide by 250 feet long.

Apply Miller's WEEDER uniformly over soil surface within 2 to 3 days following cultivation to prevent sub-surface germinating weeds from becoming established before the chemical is activated.

If there is no effective rain within 3 to 4 days after application, sprinkle the area treated to secure best results. The less the plant bed is disturbed after application, the more effective the application will be. Weed control up to four months or more may be expected following proper application.

Where possible application should be made in early spring. Applications later in the season can be made provided beds have been cultivated and are free of weeds.

WHERE TO APPLY

VEGETABLE GARDENS

General Directions—make applications as noted under "How to Apply" or as noted under specific crop.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE—Apply at seeding or transplanting. Granules can be applied directly over transplants without injury.

BEANS (Snap or Dry)—Apply at seeding.

CANTALOUPE, CUCUMBERS, HONEYDEW MELONS, SQUASH (summer and winter) WATERMELONS—Apply 4 to 6 weeks after seeding.

COLLARDS, KALE, MUSTARD GREENS, TURNIPS—Apply at seeding.

EGGPLANTS, PEPPERS—Apply at transplanting. Granules can be applied directly over transplants without injury.

GARLIC, ONIONS—Apply at seeding or transplanting. Granules can be applied directly over transplants without injury.

LETTUCE, LEAF—Make one application between 1 week and no later than 3 weeks after emergence.

LETTUCE, HEAD—Make one application only between 1 to 5 weeks after seeding.

POTATOES (whole or cut pieces)—Apply at planting.

SWEET POTATOES, YAMS—Apply at transplanting.

TOMATOES—Apply 4 to 6 weeks after transplanting. Granules can be applied directly over transplants without injury.

CAUTION

Keep out of reach of children. Do not take internally. Avoid contact with eyes. Wash after handling or using.

Do not store near foodstuffs and feed, seed, fertilizers or other pesticides in order to prevent contamination. Store in a dry place.

Use only on vegetables, flowers and shrubs listed.

Do not graze livestock on treated areas or feed plant refuse to livestock.

Destroy empty container. Do not re-use for any purpose.

CAUTION Keep out of the reach of children
See right side panel for additional cautions