

Use only as directed, noting all Cautions and Warnings.

GENERAL DIRECTIONS

1. Dust lightly on first appearance of insect or insect injury.
2. Make complete coverage of all pests and all parts of plants especially upper and lower leaf surfaces.
3. Repeat applications as necessary for control, but observe the number of days before harvest for last application noted by each crop (for instance on Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi you may apply this dust up to 3 days before picking or harvest).
4. Repeat applications if rain occurs within 24 hours after application of dust.

VEGETABLES

BEANS (day of harvest)—for control of Flea Beetles, Mexican Bean Beetles, Leafhoppers, Cucumber Beetles, Western Bean Cutworm.

CABBAGE, BRUSSELS SPROUTS, BROCCOLI, CAULIFLOWER, KOHLRABI (3)—for control of Flea Beetles, Armyworms, Cabbage Caterpillars, Corn Earworm.

CORN (day of harvest) — for control of Corn Earworm, Corn Rootworm Adults, Fall Armyworms, Leafhoppers, Flea Beetles. For **Corn Earworm**, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals.

CUCUMBERS, MELONS, PUMPKINS, SQUASH (Summer and Winter) (day of harvest) — for control of Cucumber Beetles, Flea Beetles, Leafhoppers, Melon Worms, Pickle Worms, Squash Bugs.

BEETS*, PARSNIPS, RADISHES, RUTABAGAS, HEAD LETTUCE (3*)—for control of Armyworms and Flea Beetles. *On Beets, do not use within 14 days of harvest if tops are to be used as food or feed.

ENDIVE, LEAF LETTUCE, MUSTARD GREENS, SPINACH (14) — for control of Armyworms, Cabbage Caterpillars, Flea Beetles, Leafhoppers, Lygus Bugs, Stink Bugs.

TOMATOES, EGGPLANTS, PEPPERS (day of harvest)—for control of Flea Beetles, Leafhoppers, Fall Armyworms, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs, Stink Bugs.

POTATOES (day of harvest)—for control of Colorado Potato Beetle, Flea Beetles, Leafhoppers.



SEVIN*

5% DUST INSECTICIDE

for
VEGETABLES — BERRIES
DOGS — CATS

For the **VEGETABLE and BERRY GARDEN**—to control Tomato Fruitworm, Flea Beetles, Spittlebugs, Orange Tortrix, Corn Earworm, Cutworms, Armyworms, Leafhoppers, and certain other garden insects.

For **DOGS and CATS** — to control Fleas and brown Dog Ticks

Active ingredient
 Carbaryl (1-naphthyl N-methylcarbamate) 5.0%
 Inert ingredients 95.0%
 U.S. Pat. Nos. 2,903,478; 3,009,855
 *SEVIN is the Reg. T.M. of the Union Carbide Corp.

CAUTION — Keep out of reach of children

See right side panel for additional cautions

NET WT. 1 LB.



BERRIES

STRAWBERRIES (1)—for control of Meadow Spittlebugs, Strawberry Leafroller and Strawberry Weevil.

RASPBERRIES, BOYSENBERRIES, BLACK-BERRIES (7) — for control of Japanese Beetles, Leafrollers, Leafhoppers, Orange Tortrix, Raspberry Aphid.

BLUEBERRIES (day of harvest)—for control of Blueberry Maggot, Cranberry Fruitworm, Orange Tortrix.

GRAPES (day of harvest) — for control of Grape Leafhoppers, Grape Leaf Skeletonizer, Red-banded Leafroller, June Beetles.

DOGS and CATS

On the animals—for control of Fleas and Brown Dog Ticks—dust liberally over the animals, avoiding the eyes, and rub thoroughly through the hair onto the skin. Repeat at weekly intervals if necessary. Do not use on kittens or puppies under 4 weeks old.

In the animal quarters—for control of Fleas and Brown Dog Ticks—dust liberally in and around dog houses, kennels, sleeping areas. Force dust into all cracks and crevices. Repeat as necessary.

CAUTION

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally.

Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating.

Note to Physician—Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of feed or food. Wash all vegetables and fruits before eating or canning.

For protection of honey bees, avoid use during periods when they are visiting the garden. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after applications.

Destroy empty container. Do not re-use for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S.D.A. Reg. No. 802-442 MP-1D

802-442
 1-29-1978
 1/2

2/2

ACCEPTED
JAN 29 1970
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 802-449

802-202F