

Please read the entire label.
Use only as directed, noting all
Cautions and Warnings.

GENERAL DIRECTIONS

- 1 Dust lightly on first appearance of insect or insect injury
- 2 Make complete coverage of all pests and all parts of plants especially upper and lower leaf surfaces
- 3 Repeat applications as necessary for control, but observe the number of days before harvest for last application noted by each crop (for instance on Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi, you may apply this dust up to 3 days before picking or harvest)
- 4 Repeat applications if rain occurs within 24 hours after application of dust

VEGETABLES

BEANS (day of harvest) — for control of Flea Beetles, Mexican Bean Beetles, Leafhoppers, Cucumber Beetles, Western Bean Cutworm

CABBAGE, BRUSSELS SPROUTS, BROCCOLI, CAULIFLOWER, KOHLRABI (3) — for control of Flea Beetles, Armyworms, Corn Earworm

CORN (day of harvest) — for control of Corn Earworms, Corn Rootworm (Adults, Fall Armyworms, Leafhoppers, Flea Beetles

For **Corn Earworm**, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce the populations

CUCUMBERS, MELONS, PUMPKINS, SQUASH (Summer and Winter) (day of harvest) — for control of Cucumber Beetles, Flea Beetles, Leafhoppers, Melon Worms, Pickle Worms, Squash Bugs

BEETS*, PARSNIPS, RADISHES, RUTABAGA, HEAD LETTUCE (3*) — for control of Armyworms and Flea Beetles. *On Beets do not use within 14 days of harvest if they are to be used as food for feed

ENDIVE, LEAF LETTUCE, MUSTARD GREENS, SPINACH (14) — for control of Armyworms, Flea Beetles, Leafhoppers, Lygus Bugs, Stink Bugs

TOMATOES, EGGPLANTS, PEPPERS (day of harvest) — for control of Flea Beetles, Leafhoppers, Fall Armyworms, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs, Stink Bugs

POTATOES (day of harvest) — for control of Colorado Potato Beetle, Flea Beetles, Leafhoppers

NET WT.
1 LB.



Sevin*

5% Dust

For the VEGETABLE and BERRY GARDEN—to control Tomato Fruitworm, Flea Beetles, Spittlebugs, Orange Tortrix, Corn Earworm, Cutworms, Armyworms, Leafhoppers, and certain other garden insects.

For DOGS and CATS—to control Fleas and brown Dog Ticks

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl
N-methylcarbamate) 5.0%
INERT INGREDIENTS 95.0%
U.S. Pat. No. 2,903,478; 3,009,856

*SEVIN is the Reg. TM of the
Union Carbide Corp.

CAUTION Keep out of reach of children.
See right side panel for additional cautions.
E.P.A. Reg No 802-442-AA