

Daria Mills
Registration Specialist
The Chas. H. Lilly Company
P.O. Box 83179
Portland, OR 97283

MAR 6 1997

Subject: Miller Sevin 5% Dust Insecticide
EPA Registration No. 802-442
Amendments dated May 3, 1996 and October 17, 1996
Deletion of Pet uses and Response to Agency letter dated 10/1/96

Dear Ms. Mills:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:
 - a) In the Directions for Use on CORN, change "2 to 3 day intervals" to "7 day intervals."
2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

/s/

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

BEST AVAILABLE COPY

2 9 4

Millers Sevin 5% Dust Insecticide

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS 95.0%

SEVIN is the Reg. T.M. of Rhone-Poulenc for carbaryl insecticide.

In the garden to control Tomato Fruitworm, Flea Beetles, Spittlebugs, Corn Earworm, Cutworms, Armyworms, Leafhoppers

Keep out of reach of children

CAUTION

See right side panel for additional cautions.

NET WEIGHT 1 LB. (454 g)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 6 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

802-442

96 OCT 23 11:15

REC'D EPA/OPP/DPD1

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

NOTE TO PHYSICIAN: Carbaryl Insecticide is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of feed or food. Wash all vegetables and fruits before eating or canning.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Please read the entire label. Use only as directed.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household use only.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin.

Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

For plants above chest height, consider using a liquid carbaryl product.

Do not repeat application more than once a week.

Dust lightly on first appearance of insect or insect injury.

Make complete coverage of all pests and all parts of plants especially upper and lower leaf surfaces.

Observe the number of days before harvest for last application noted by each crop (for instance on cabbage, broccoli, brussels sprouts, cauliflower and kohlrabi, you may apply this dust up to 3 days before picking or harvest).

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

VEGETABLES

BEANS (day of harvest) - for control of Flea Beetle, Mexican Bean Beetles, Leafhoppers, Cucumber Beetles, Western Bean Cutworm.

CABBAGE, BRUSSELS SPROUTS, BROCCOLI, CAULIFLOWER, KOHLRABI (3) - for control of Flea Beetles, Armyworms, Corn Earworm.

CORN (day of harvest) - for control of Corn Earworm, Corn Rootworm Adults, Fall Armyworms, Leafhoppers, Flea Beetles.

For Corn Earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

CUCUMBERS, MELONS, PUMPKINS, SQUASH (Summer and Winter) (day of harvest) - for control of Cucumber Beetles, Flea Beetles, Leafhoppers, Melonworms, Pickleworms, Squash Bugs.

BEETS*, PARSNIPS, RADISHES, RUTABAGAS, HEAD LETTUCE (3*) - for control of Armyworms and Flea Beetles. *On Beets, do not use within 14 days of harvest if tops are to be used as food or feed.

ENDIVE, LEAF LETTUCE, MUSTARD GREENS, SPINACH (14) - for control of Armyworms, Flea Beetles, Leafhoppers, Lygus Bugs, Stink Bugs.

TOMATOES, EGGPLANTS, PEPPERS (day of harvest) - for control of Flea Beetles, Leafhoppers, Fall Armyworms, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs, Stink Bugs.

POTATOES (day of harvest) - for control of Colorado Potato Beetle, Flea Beetles, Leafhoppers.

BERRIES

STRAWBERRIES (1) - for control of Meadow Spittlebugs, Strawberry Leafroller and Strawberry Weevil.

RASPBERRIES, BOYSENBERRIES, BLACKBERRIES (7) - for control of Japanese Beetles, Leafrollers, Leafhoppers, Orange Tortrix, Raspberry Aphid.

BLUEBERRIES (day of harvest) - for control of Blueberry Maggot, Cranberry Fruitworm, Orange Tortrix.

GRAPES (7) - for control of Grape Leafhoppers, Grape Leaf Skeletonizer, Red-Banded Leafroller, June Beetles.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals.

Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of the product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Filled by weight not by volume. Product may settle after packaging.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97203
EPA Reg. No. 802-442

EPA Est. 44616-MO-01

77652