#### OUTSIDE OF HOMES

#### ANTS, EARWIGS, SPIDERS, BOX ELDER BUGS

- 1. Use 1 tablespoon per 1 gal. of water (1 pt. per 32 gal.).
- 2. Spray outside areas thoroughly. Apply around house foundations, garbage containers; along runways; in cracks of cement walks and patios. Also spray Box Elder trees and other ornamentals where bugs congregate, using 1 teaspoon per ] gal, of water.
- 3. Repeat as necessary.

#### **TERMITES** in Buildings

- 1. Use 1 pt. mixed with water to make 12 gallons of mixed Spray.
- 2. Homes with Basements or Crawl Spaces, New Construction: Dig trench along inside and outside of foundations 15 to 18 inches deep, but not below footing, (8 to 12 in. wide). Use 2 gal, of mixed spray for each 5 linear feet, applying half the amount to the bottom of trench against foundation. Apply remainder of mixed spray proportionately at 6 inch intervals while back filling trench, making last application just below ground level and covering with a thin layer of soil. For trenches 18 to 30 inches deep but not below footing use 4 gal. of mixed spray for each 5 linear feet.
- 3. Slab type Construction—Apply 1 gal. mixed spray per 10 sq. ft. of soil surface as an over-all preconstruction treatment under slab for porches, entrance platforms and utility entrances.
- 4. Inspect area yearly to determine possible reinfestation, and re-treat when necessary.

#### WARNING

Harmful if swallowed. Avoid breathing spray mist. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water.

Do not spray into or near fire or open flame. Do not smoke while spraying. Avoid contamination of feed and foodstuffs. Do not contaminate drinking water.

Do not use this product on forage or feed crops which may be fed to dairy animals or animals being finished for slaughter In order to avoid injury to bees, do not apply to plants in bloom.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

. .

.

- ;

Destroy empty container. Do not re-use for any purpose.

\*\*\* • • • • • •



NET CONTENTS

ONE PINT

#### ACTIVE INGREDIENTS

\*Chlordane, technical .... 72.3% Petroleum Distillate .... 21.5% INERT INGREDIENTS ..... 6.2% \*Equivalent to 43.38% octachloro-4, 7-methanotetrahydroindane and

28.92% related compounds.

WARNING Keep out of the reach of children See left side panel for additional warnings

E.P.A. Reg No. 802-443-AA

#### Please read entire label. Use only as directed, noting all cautions.

\$**3**79

#### LAWNS

## SOD WEBWORM (Lawn Moth), WHITE GRUBS, JAPANESE BEETLE LARVAE, ANTS, CUTWORMS, MOLE CRICKETS, ARMYWORMS

- 1. Use 1 tablespoon per 1 gal. of water (1 pt. per 32 gal.).
- 2. Apply evenly over the lawn at the rate of 1 gal. per 150 sq. ft.
- 3. To control Nighterawlers, use 4 oz. in 10 to 15 gal. of water. Apply evenly over TOOO sq. ft. of lawn.
- 4. For best results apply in the evening when lawn is dry. After treatment, follow with moderate sprinkling  $(\frac{1}{4} \text{ to } \frac{1}{2} \text{ inch})$ of water) to thoroughly wet the soil.
- 5. Keep children or pets off treated areas until insecticide is washed into soil and grass is thoroughly dry

### ORNAMENTAL GARDENS

STRAWBERRY ROOT WEEVIL (Brachyrhinus) ( CUTWORMS, BLACK VINE WEEVIL — Rhododendrons, Azaleas, Camellias, Primroses

- 1. Use 1 teaspoon per 1 gal. of water (1 pt. per 100 gal.).
- 2. Established Plantings-apply generously around the base of plants, Repeat as necessary.
- 3. New Seed Beds—apply at the rate of 1 gal. diluted material per 40 sq. ft. soil area and work thoroughly into top 4 to 6 inches of soil before planting.

THRIPS, ANTS, DIABROTICA BEETLES, FLEA BEETLES, LEAF MINERS, LEAFROLLERS, SOWBUGS, LYGUS BUGS, STINK BUGS, EARWIGS, GRASSHOPPERS on Roses, Gladiolus, Chrysanthemums Dahlies, Asters, Petunias, Carnations, Marigolds, Pansies, Zinnias.

- 1. Use 1 teaspoon per 1 gal. of water.
- 2. Spray plants thoroughly to cover both top and undersides of leaves as well as soil around plants. Repeat as necessary.

NOTICE: Follow directions carefully. Timing and method of application. weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewitn.

MP-6J

THE CHAS. H. LILLY CO. PORTLAND, OREGON EPA Est. 802-OR-1

#### OUTSIDE OF HOMES

#### ANTS, EARWIGS, SPIDERS, BOX ELDER BUGS

- <sup>1</sup> Use 1 tablespoon per 1 gal of water (1 pt per -2 gal)
- 2. Spray outside areas thoroughly. Apply around house foundations garbage containers, along runiways, in cracks of ce ment walks and patios. Also spray Box Elder frees and other ornamentals where bugs congregate, using 1 teaspeen per I gal, of water
- Repeat as necessary

#### **TERMITES** in Buildings

- 1. Use 1 pt. mixed with water to make 12 gallons of mixed Spray.
- 2 Homes with Basements or Crawl Spaces, New Construction: Dig trench along inside and outside of foundations 15 to 15 inches deep, but not below footing, 18 to 12 in. wider. Use 2 gal. of mixed spray for each 5 linear feet, applying half the amount to the bottom of trench against foundation Apply remainder of mixed spray proportionately at 6 inch intervals while back filling trench, making last application just below ground level and covering with a thin layer of soil For trenches 18 to 30 inches deep but not below footing use 4 gal. of mixed spray for each 5 linear feet.
- Slab type Construction—Apply 1 gal, mixed spray per 10 sq ft. of soil surface as an over-all preconstruction treatment under slab for porches, entrance platforms and utility entrances.
- Inspect area yearly to determine possible reinfestation, and re-treat when necessary.

#### WARNING

Harmful if swallowed. Avoid breathing spray mist, Contact with skin can cause toxic symptoms. In case of contact with Kin, wash with soap and water.

Do not spray into or near fire or open flame. Do not smoke while spraying. Avoid contamination of feed and foodstuffs, Do not contaminate drinking water.

Do not use this product on forage or feed crops which may be fed to dairy animals or animals being finished for slaughter In order to avoid injury to bees, do not apply to plants in bloom.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label

Destrivempty container. Do not re-use for any purplise

# NET CONTENTS ONE PINT



# 800

ACTIVE INGREDIENTS

\*Chlordane, technical Petroleum Distillate

72.3°5 21.5°, **INERT INGREDIENTS** 6.2% \*Equivalent to 43.38% octachloro-4. 7-methanotetrahydroindane and 28.92% related compounds.

WARNING Keep out of the reach of children See left side panel for additional warnings

- No. 802-443-AA E.P.A

Please read entire label. Use only as directed, noting all cautions.

#### LAWNS

#### SOD WEBWORM (Lawn Moth), WHITE GRUBS, JAPANESE BEETLE LARVAE, ANTS, CUTWORMS, MOLE CRICKETS, ARMYWORMS

- 1. Use 1 tablespoon per 1 gal of water (1 pt. per 32 gal.)
- 2 Apply evenly over the lawn at the rate of 1 gal, per 150 sg ft.
- 3 To control Nighterawlers, use 4 oz. in 10 to 15 gal of water Apply evenly over 1000 sg. ft. of lawn.
- 4. For best results apply in the evening when lawn is dry. After treatment, follow with moderate sprinkling (14 to 12 inch of water: to thoroughly wet the soil.
- 5 Keep children or pets off treated areas until in chicide is washed into soil and grass is thoroughly dry

#### ORNAMENTAL GARDENS

STRAWBERRY ROOT WEEVIL (Brachyrhinus) ( CUTWORMS, BLACK VINE WEEVIL - Rhododendrons, Azaleas, Camellias, Primroses

- 1. Use 1 teaspoon per 1 gal, of water (1 pt. per 100 gal.).
- 2 Established Plantings—apply generously around the base of plants. Repeat as necessary,
- 5. New Seed Beds apply at the rate of 1 gal. diluted material per 40 sq. ft. soil area and work thoroughly into top 4 to 6 inches of soil before planting.

THRIPS, ANTS, DIABROTICA BEETLES, FLEA BEETLES, LEAF MINERS, LEAFROLLERS, SOWBUGS, LYGUS BUGS, STINK BUGS, EARWIGS, GRASSHOPPERS on Roses, Gladiolus, Chrysanthemums Dahlies, Asters, Petunias, Carnations, Marigolds Pansies, Zinnias

- 1. Use 1 teaspoon per 1 gal of water
- 2. Spray plants thoroughly to cover both tup and undersides of leaves as well as soil around plants. Repeat as necessary

NOTICE: Follow directions carefully. Timing and method of application. weather and crep conditions, mixtures with other chemicals not specifically recommended, and other influencing rators in the use of this product are heyond the control of the scher. Buyer assumes all risks of use, storage or handling of this inaterial not in strict accordance with directions given. he rewith:

MP GJ

THE CHAS. H. LILLY CO. PORTLAND, OREGON EPA Est 802-OR-1

\$**3**79

