



# 7.5% DITHANE\* M-45 DUST

### ACTIVE INGREDIENT:

A coordination product of zinc ion and manganese ethylene bisdithiocarbamate in which the ingredients are: ..... 6.00%

Manganese ..... 1.20%

Zinc ..... 0.15%

Ethylene bisdithiocarbamate ..... 4.65%

INERT INGREDIENTS ..... 94.00%

TOTAL ..... 100.00%

EPA Reg. No. 802-437

Keep out of reach of children

## CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May cause irritation of nose, throat eyes and skin. Do not breathe dust. Protective clothing such as long-sleeve shirts, long pants, rubber gloves, and rubber boots are to be worn by persons involved during mixing and loading of this product. In cases of skin contact, wash with water; for eyes, wash copiously with water and get medical attention.

#### ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NET WT. 50 LBS.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

**Important:** Keep away from fire and sparks. Store in a cool dry place. Keep container closed when not in use.

THE CHAS. H. LILLY CO  
Portland, Oregon 97218  
EPA Est. 802-OR-1

### SUGGESTED USE

#### POTATO SEED PIECE TREATMENT FOR PROTECTION AGAINST FUSARIUM SAMBUCINUM

Lilly/Miller 7.5% DITHANE M-45 DUST is a special fungicide formulation for application on whole or cut potato tubers to protect them from seed-piece decay caused by *Fusarium sambucinum*.

Use 1 lb. of dust per 100 lbs. of whole or cut potato seed pieces. Apply the dust in a manner such that the entire surface of the seed-piece becomes thoroughly covered with the fungicide dust. Plant seed as soon after treatment as possible.

**DO NOT USE TREATED SEED PIECES FOR FOOD OR FEED PURPOSES.**

\*DITHANE is the Reg. T.M. of the Rohm and Haas Co.

**WARRANTY:** The use of this material being subject to conditions beyond our knowledge and control, the seller makes no warranty of any kind, express or implied, concerning the effects of this product, when not used in accordance with directions.

LC-3S

BEST DOCUMENT AVAILABLE

ACCEPTED  
AUG 31 1983  
Federal Insecticide, Fungicide, and Rodenticide Act  
EPA-700-1-83-113/