

MADE
OCT. 10, 1972
802-464

10/10/72
MADE IN U.S.A.

DIRECTIONS FOR USE

WHITE PINE WEEVIL—on Norway Spruce, and White, Jack, Red, Limber, Foxtail and Western White Pines

1. Use 2 quarts per 6 gallons of water.
2. Apply to the upper half of each leader of trees during late March or early April.

STRAWBERRY ROOT WEEVILS—On Ornamentals, Blackberries, Blueberries, Boysenberries, Dewberries, Raspberries.

1. Use 2 gallons per acre in sufficient water to obtain uniform coverage.
2. Mix thoroughly in the top 6 inches of soil before planting.
3. Make no application while edible parts are present.

WIREWORMS—Corn, Tomatoes, Peppers

1. Use 1 to 1 1/2 gallons per acre in sufficient water to obtain uniform coverage.
2. Make application before discing or rototilling and mix thoroughly with top 6 to 8 inches of soil.
3. Do not apply after edible parts start to form.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MI 76



HEPTACHLOR 2E

(2 lbs. Heptachlor per Gallon)

**WHITE PINE WEEVIL
STRAWBERRY ROOT WEEVILS
NARCISSUS BULB FLY • WIREWORMS**

Active ingredients	
*Heptachlor	23.4%
Related Compounds	7.8%
Aromatic Petroleum Derivative Solvents	64.7%
Inert ingredients	4.1%
*Heptachlorotetra-hydro-4,7-methanoindene	

E.P.A. Reg. No. 802-464-AA

WARNING — Keep out of reach of children

See right side panel for additional warnings

_____ GALLON

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214

MIXING DIRECTIONS

Add half the required amount of water to the sprayer, then add the specified amount of Miller's HEPTACHLOR 2E, with agitation, then add balance of the water with continued agitation.

DIRECTIONS FOR USE

NARCISSUS BULB FLY—Narcissi

1. Use 1 quart in 50 gallons of water (4 teaspoons in 1 gallon).
2. Dip bulbs for 10 minutes prior to planting.
3. After bloom, when fly appears, drench base of stem using above dilution and applying with a sprinkling can.

WARNING

May be fatal if swallowed. Do not breathe vapor or spray mist. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed or foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by breaking or by punching holes in them and burying with wastes.