

# METHOXYCHLOR 2E

(2 lbs. Methoxychlor per gallon)

## DIRECTIONS FOR USE

**ALFALFA, CLOVER . . . Seed or Hay . . .** Alfalfa Weevils, Spittle Bugs, Leafhoppers

1. Use 2 to 3 quarts per acre. Apply in sufficient water to give good coverage.
2. Begin applications at first sign of insects and repeat as infestations warrant, but do not apply within 7 days of harvest or grazing.

**BEANS . . .** Mexican Bean Beetles, Leafhoppers

1. Use 2 quarts per acre. Apply in sufficient water to give good coverage of both upper and lower leaf surfaces.
2. Repeat in 2 weeks if infestations warrant, but do not apply within 3 days of harvest.

**PEAS . . .** Pea Leaf Weevil, Pea Weevil

1. Apply at the rate of 3 qts. per acre. Use sufficient water to insure thorough coverage (5 to 10 gal. by air; 20 gal. by ground application). Repeat applications as needed to maintain control. Do not apply within 7 days of harvest.
2. For the control of **Pea Weevil**, apply at the above rates during early bloom period before eggs are laid. Repeat in 5 days.

**SOYBEANS, COWPEAS . . .** Spittle Bugs, Leafhoppers

1. Use 2 quarts per acre. Apply in sufficient water to give good coverage.
2. Treat at first sign of insects and repeat as infestations warrant, but do not apply within 7 days of harvest, or grazing.

**STORED BARLEY, CORN, OATS, RYE, WHEAT . . .** Saw Toothed Beetles, Flat Grain Beetles, Granary Weevils

1. Use ½ cup (8 tablespoons) in one gallon of water, and apply to each 500 sq. ft. of grain surface. Make no more than 2 applications.

## CAUTION

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin. Wash thoroughly after using.

Avoid contamination of feed and foodstuffs. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

Highly toxic to fish. Do not contaminate lakes, ponds and streams. Do not apply where runoff is likely to occur. Toxic to bees and should not be applied when bees are actively visiting the area.

Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment or disposal of wastes

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

## LIVESTOCK SPRAYS

**CAUTION: Do not use on Dairy Animals**

**HORN FLIES (Beef Cattle)**

1. As a direct application, use 1 gallon per 50 gallons of water (4/5 pt. per 5 gallons of water).
2. Spray 2 quarts of the mixture per animal.
3. Apply by means of a power sprayer with orchard type nozzle or knapsack sprayer, wetting shoulders, back and flanks.
4. Repeat every 3 to 4 weeks during fly season.

**BACK RUBBER TREATMENT . . .** Hornflies on Beef Cattle; and as an aid in the reduction of Lice on Beef Cattle; Lice on Hogs

1. Mix 1 quart with 4 quarts stove oil or light grade of fuel oil.
2. Pour 1 gallon of this mixture evenly over 15 to 20 feet of burlap "cable" in each of several rubbing units installed in areas where the animals loaf or feed.
3. The burlap should be retreated every 3 to 5 weeks using 2 quarts of the mixture per 15 to 20 feet of "cable."
4. Regular exposure of the animals for about 10 weeks is necessary for best results.

**LICE on Beef Cattle and Hogs**

1. Direct spraying . . . use 1 quart per 10 to 12 gallons of water.
2. Spray so as to insure thorough coverage of animals, using one quart of spray mixture per hog and 2 quarts of spray per beef animal.

- 5 GALLONS OF WATER).
2. Spray 2 quarts of the mixture per animal.
  3. Apply by means of a power sprayer with orchard type nozzle or knapsack sprayer, wetting shoulders, back and flanks.
  4. Repeat every 3 to 4 weeks during fly season.

**BLACK RUBBER TREATMENT . . .** Horns on Beef Cattle; and as an aid in the reduction of Lice on Beef Cattle; Lice on Hogs

1. Mix 1 quart with 4 quarts stove oil or light grade of fuel oil.
2. Pour 1 gallon of this mixture evenly over 15 to 20 feet of burlap "cable" in each of several rubbing units installed in areas where the animals loaf or feed.
3. The burlap should be retreated every 3 to 5 weeks using 2 quarts of the mixture per 15 to 20 feet of "cable."
4. Regular exposure of the animals for about 10 weeks is necessary for best results.

**USE on Beef Cattle and Hogs**

1. Direct spraying . . . use 1 quart per 10 to 12 gallons of water.
2. Spray so as to insure thorough coverage of animals, using one quart of spray mixture per hog and 2 quarts of spray per beef animal.

# METHOXYCHLOR 2E

(2 lbs. Methoxychlor per gallon)

## AGRICULTURAL INSECTICIDE

for use on

**Alfalfa, Clover, Beans, Peas, Soybeans, Cowpeas  
and Stored Grain  
Beef Cattle and Hogs**

PROTECT FROM TEMPERATURES BELOW 32° F.  
FLAMMABLE MIXTURE . . . Keep away from heat or open flame

ACTIVE INGREDIENTS	
Technical Methoxychlor*	25.0%
Aromatic Petroleum Derivative Solvents	66.2%
INERT INGREDIENTS	8.8%
Total	100.0%

\*Equivalent to 22% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 3% other isomers and reaction products

**CAUTION** — Keep out of reach of children  
See back panel for additional cautions

E.P.A. Reg. No. 802-466-AA

**NET CONTENTS \_\_\_\_\_ GALLON(S)**

1. Use 1 gallon per 10 gallons of water.
2. Apply at the rate of 1 gallon of finished spray mixture per 500 sq. ft. of surface.
3. Spray particularly those areas where flies crawl or rest and congregated.
4. Repeat applications as infestation warrant.
5. Protect from the spray application all drinking utensils, feed buckets, milking utensils, milk and feed troughs.
6. All animals should be excluded from barns until after the spray application has been completed.

**RESIDUAL SPRAY in Flour Mills, Grain and other Cereal Storages . . .** Toothed Beetles, Flat Grain Beetles, and many Weevils

1. Use 1 gallon per 10 gallons of water (2 qts. per 5 gal.), as a residual application to woodwork in bins and other grain storages which previously have been thoroughly cleaned.
2. Apply by means of a knapsack garden or power sprayer at the rate of 2 gallons per 1000 sq. ft. for control of the above insects which are exposed or become accessible after emerging from burrows and crevices of the woodwork after cleaning.