

HEPTACHLOR 2.5G

(Granular Product Containing 2.5% Heptachlor)

Active ingredients	
Heptachlor (Heptachlorotetrahydro 4,7-methanoindene)	2.5%
Related Compounds	0.9%
Inert ingredients	96.6%

CAUTION — Keep out of reach of children

Harmful if swallowed. Avoid inhalation of dust. Avoid skin contact and wash hands, arms and face with soap and water after using.

USE PRECAUTIONS

Read entire label. Use strictly in accordance with directions, cautions and warnings on label and in conformity with Federal and State Regulations.

Avoid contamination of feed and foodstuffs.

Apply this product only as specified on this label. Do not apply when weather conditions favor drift of any dust from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is liable to occur.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

800 417
 Jul 14, 1972
 SUGGESTED USES
 SOIL APPLICATIONS ONLY

Crops and Pests	Pounds per Acre
<u>CABBAGE</u> - Cabbage Maggots	30 to 40
Apply to soil surface around the plants immediately after transplanting.	
<u>LETTUCE</u> - Wireworms	80 to 120
Preplanting soil application only. Apply to soil surface and mix with top 3 to 6 inches of soil before planting.	
<u>RUTABAGAS</u> - Root Maggots	80 to 120
Apply to soil soon after plants emerge with a second applications 3 to 4 weeks later. Do not use tops as food or feed.	
<u>TOMATOES, PEPPERS</u> - Wireworms	80 to 120
Apply to soil surface and mix with top 3 to 6 inches of soil before planting. Do not apply after edible parts start to form.	
<u>ORNAMENTAL NURSERY STOCK AND ORNAMENTALS</u>	
<u>Root Weevils</u> (Brachyrhinus)	200
Apply to soil prior to or at time of setting plants in field. Work into top 3 to 6 inches of soil as soon as possible after transplanting.	