


[802-473]

"SAVIN" copy of printed label --



802-473  
ACCEPTED  
NOV 27 1972  
FEDERAL BUREAU OF INVESTIGATION  
U.S. DEPARTMENT OF JUSTICE  
SUBJECT TO ATTACHED COMMENTS.

**DIRECTIONS FOR USE**

**SPRING AND SUMMER SPRAY**

1. Use 3 tablespoons per gallon of water.
2. Apply as a thorough wetting spray as insects appear or before disease becomes evident.
3. Repeat applications at 7 to 10 day intervals as needed to maintain control. (See Caution.)

Use on ROSES, FLOWERS, SHRUBS to control: Aphids, Flea Beetles, Leafhoppers, Leaf Tiers, Thrips (exposed), Powdery Mildew, Black Spot.

Use on BEANS, PEAS for control of: Mexican Bean Beetle, Leafhoppers, Flea Beetles, Pea Weevil, Aphids. For control of Pea Weevil, spray foliage during early bloom period before eggs are laid.

Use on BROCCOLI, BRUSSELS SPROUTS, CABBAGE, KALE, KOHLRABI, LETTUCE, CELERY for control of: Aphids (other than Woolly), Imported Cabbage Worms, Diamondback Moth Larvae, Flea Beetles, Tarnished Plant Bugs, Vegetable Weevils, Celery Looper.

Use on CANTALOUPE, CUCUMBERS, MELONS, PUMPKINS AND SQUASH (Summer and Winter) for control of: Melon Aphids, Leafhoppers, Squash Bug Nymphs, Spotted and Striped Cucumber Beetles.

Use on PEPPERS for control of: Aphids.

Use on TOMATOES for control of: Blister Beetles, Aphids, Flea Beetles, Colorado Potato Beetle.

Use on RASPBERRIES for control of: Rednecked Cane Borer, Raspberry Cane-Borer, Raspberry Fruitworm, Thrips (exposed), Leafhoppers. For Raspberry Cane Borer and Rednecked Cane Borer, spray foliage just before plants bloom. Repeat in about 2 weeks. For Raspberry Fruitworm, spray foliage 7 days after first blossom. Repeat in 10 days and again 10 days later.

Use on STRAWBERRIES for control of: Spittlebugs.

Use on CHERRIES for control of: Cherry Fruit Fly. Thorough full coverage applications should be made beginning when adult flies appear, usually just after the blossom buds open. A second application should be made after the shucks have fallen. Repeat every 10 days, or as required, to within 1 day of harvest.

Use on GRAPEFRUIT, LEMONS, ORANGES for control of: Citrus Aphids.

ONE QUART

\$4.49



**CAUTION** Keep out of the reach of children  
See right side of panel for additional cautions.  
E. P. A. Reg. No. 802-473

**DIRECTIONS FOR USE**

**LAWNS AND ORNAMENTAL TURF**—Lawn Moth, Sod Webworm

1. Use 3 tablespoons per gallon of water.
2. Sprinkle or spray one gallon of mixture on approximately 20 sq. ft. of lawn or turf.

**DORMANT AND DELAYED DORMANT CLEANUP SPRAY**

1. Use 6 tablespoons per gallon of water.
2. Apply carefully to insure that the entire tree is thoroughly wet.

Use on PEACHES to control: Peach Leaf Curl. Apply in late fall or early winter before fall rains and again before buds swell.

Use on APPLES, PEARS to control: Powdery Mildew, Scab. Apply in late winter and again just before blossoming.

Use on APRICOTS, PRUNES, PLUMS to control: Brown Rot Blossom Blight. Apply in late winter and repeat at petal fall.

**CAUTION**

Harmful if swallowed. Avoid contact with skin. Avoid breathing of spray mist.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not use on Peas within 7 days of harvest, or on other food crops named on this label within 1 day of harvest.

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Destroy empty container. Do not reuse for any purpose.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-1G

— MILLER PRODUCTS —  
THE CHAS. H. LILLY CO.  
PORTLAND, ORE. 97214