

RECEIVED  
12 AUG 1974  
REGISTRATION DIVISION  
PESTICIDES, EPA

NET WT.  
1 lb.

Miller's

**VEGETABLE DUST**

**CAUTION**  
Harmful if swallowed or inhaled. May cause irritation of nose and throat. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after using and before eating or smoking.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Wash all edible portions of treated plants thoroughly before eating or processing. Avoid storage of this product near food and food products. Do not use treated tops of vegetables for food or feed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Destroy empty container. Do not reuse for any purpose.

**ACTIVE INGREDIENTS:**

Carbaryl (1-Naphthyl N-Methylcarbamate)	5.0%
Rotenone	1.0%
Other Cube Resins	1.0%
Zinc (Zinc Ethylene Bisdithiocarbamate)	5.0%
<b>INERT INGREDIENTS</b>	<b>88.0%</b>

U.S. Pat. Nos. 2,103,178, 2,009,855  
SEVIN is the Reg. TM of the Union Carbide Corp.

**NOTICE:** Follow directions carefully. Timing and method of applications, weather, and crop conditions, together with other chemicals not specifically recommended, are other influencing factors in the use of this product beyond the control of the seller. Buyer assumes all use, storage or handling of this material in accordance with directions given hereon.

# Vegetable dust

**INSECTICIDE FUNGICIDE  
FOR  
TOMATOES, CARROTS, PEAS  
CORN, CABBAGE, BEANS  
MELONS AND CUCUMBERS**

**ACCEPTED**  
DEC 26 1974  
UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
ED UNL... C... POISON REGISTER  
TO UNL... C... POISON REGISTER

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON  
EPA Est. 802-OR-1

**CAUTION** Keep out of the reach of children  
See left side panel for additional cautions  
E.P.A. Reg. No. 802-474-AA

**GENERAL DIRECTIONS**

Shake duster vigorously to mix air with dust thus insuring light uniform dust applications.

**INSECT CONTROL**

1. Dust lightly on first appearance of insect or insect damage.
2. Make complete coverage of all foliage.
3. Repeat applications every 7 to 10 days as necessary for control but be certain to observe the number of days between last application and harvest as shown in parenthesis following each crop listed.

**BEANS (7)**

Cucumber Beetles, Flea Beetles, Leafhoppers, Armyworm, Stink Bugs.

**BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)**

Flea Beetles, Imported Cabbageworm, Stink Bugs, Armyworm.

**CARROTS (7)**

Flea Beetles, Leafhoppers, Spittlebugs, Stink Bugs, Lygus Bugs.

**CORN (0)**

Fall Armyworm, Flea Beetles, Leafhoppers, For Corn Earworm control see Specific Directions.

**CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)**

Cucumber Beetles, Pickleworm, Melonworm, Flea Beetles, Leafhoppers, Squash Pests.

**HEAD LETTUCE (10), LEAF LETTUCE (14)**

Flea Beetles, Leafhoppers, Armyworm, Imported Cabbageworm, Spittlebugs, Stink Bugs.

**PEAS (10)**

Leafhoppers, Armyworm, Grasshoppers.

**TOMATOES (5), PEPPERS (1), EGGPLANT (1)**

Flea Beetles, Leafhoppers, Tomato Hornworm, Tomato Fruitworm, Fall Armyworm, Stink Bugs.

**DISEASE CONTROL**

1. Dust lightly so as to insure coverage of foliage.
2. Make regular light dusting at regular intervals until disease is controlled. Observe the number of days between last application and harvest as shown in parenthesis following each crop listed.

**BEANS (7), PEAS (10)**

Anthracnose, Downy Mildew

**BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7)**

Downy Mildew, Leaf Spot

**CARROTS (7)**

Early Blight, Late Blight

**CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)**

Downy Mildew, Blossom Blight

**EGGPLANT (1)**

Early Blight, Late Blight, Cercospora Leaf Spot

**HEAD LETTUCE (10), LEAF LETTUCE (14)**

Downy Mildew, Botrytis

**PEPPERS (1)**

Cercospora Leaf Spot, Downy Mildew

**TOMATOES (5)**

Early Blight, Late Blight, Septoria Leaf Spot

**SPECIFIC DIRECTIONS**

- Corn - Corn Earworm:**
1. Dust the silk and hills.
  2. Make first application of hills show silk.
  3. Repeat at 2 to 3 total of 3 or more applications.
  4. Do not feed forage to animals before application.
  5. Applications of this product during the period will seriously reduce damage.

**Cameraman's Note**  
Poor Copy





NET WT.  
1 lb.

# Vegetable dust

**INSECTICIDE FUNGICIDE  
FOR  
TOMATOES, CARROTS, PEAS  
CORN, CABBAGE, BEANS  
MELONS AND CUCUMBERS**

**UTION**  
ved or inhaled. May  
nose and throat. Avoid  
and contact with eyes.  
Wash thoroughly after  
ing or smoking.

**ICIAN:** Carbaryl is a  
ible, cholinesterase  
antidotal.

ions of treated plants  
eating or processing  
his product near feed  
Do not use treated  
or food or feed.

ghly toxic to bees ex-  
atment or residues on  
information may be ob-  
cooperative Agricultural

ic to fish. Keep out of  
ponds. Do not apply  
y to occur. Do not ap-  
conditions favor drift.  
Do not contaminate  
ng of equipment or  
Apply this product only  
label.

ainer. Do not reuse for

NTS  
hyl  
ate) 5.0%  
1.0%  
S 1.0%  
ene  
ate) 5.0%  
TS 88.0%  
1009.855  
be

is carefully. Time of treatment  
and crop conditions, insect  
specifically recommended  
in the use of this product  
seller. Buyer assumes  
of the material  
divor here with

H. LILLY CO.  
ND, OREGON.  
802-OR-1

502-474

**ACCEPTED**  
DEC 26 1974  
UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED USE

**CAUTION** Keep out of the reach of children  
See left side panel for additional cautions  
E.P.A. Reg. No. 802-474-AA

*Miller's*

## VEGETABLE DUST

### GENERAL DIRECTIONS

Shake duster vigorously to mix air with dust thus insuring light uniform dust applications.

### INSECT CONTROL

1. Dust lightly on first appearance of insect or insect damage.
2. Make complete coverage of all foliage.
3. Repeat applications every 7 to 10 days as necessary for control but be certain to observe the number of days between last application and harvest as shown in parenthesis ( ) following each crop listed.

### BEANS (7)

Cucumber Beetles, Flea Beetles, Leafhoppers, Armyworm, Stink Bugs.

### BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)

Flea Beetles, Imported Cabbageworm, Stink Bugs, Armyworm.

### CARROTS (7)

Flea Beetles, Leafhoppers, Spittlebugs, Stink Bugs, Lycus Bugs.

### CORN (0)

Fall Armyworm, Flea Beetles, Leafhoppers, Fall Corn Earworm control see Specific Directions.

### CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)

Cucumber Beetles, Pickleworm, Melonworm, Flea Beetles, Leafhoppers, Stink Bugs.

### HEAD LETTUCE (10), LEAF LETTUCE (14)

Flea Beetles, Leafhoppers, Armyworm, Imported Cabbageworm, Spittlebugs, Stink Bugs.

### PEAS (10)

Leafhoppers, Armyworm, Grasshoppers.

### TOMATOES (5), PEPPERS (1), EGG PLANT (1)

Flea Beetles, Leafhoppers, Tomato Hornworm, Tomato Fruitworm, Fall Armyworm, Stink Bugs.

### DISEASE CONTROL

1. Dust lightly so as to make complete coverage of foliage at first appearance of disease or apply as a preventative before disease appears.
2. Make regular light applications at 7 to 10 day intervals until disease is under control but observe the number of days between last application and harvest as shown in parenthesis ( ) following each crop listed.

### BEANS (7), PEAS (10)

Anthrachnose, Downy Mildew, Rust.

### BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)

Downy Mildew, Leaf Spot.

### CARROTS (7)

Early Blight, Late Blight.

### CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)

Downy Mildew, Blossom Blight.

### EGGPLANT (1)

Early Blight, Late Blight, Anthracnose, Cercospora Leaf Spot.

### HEAD LETTUCE (10), LEAF LETTUCE (14)

Downy Mildew, Botrytis Blight.

### PEPPERS (1)

Cercospora Leaf Spot, Anthracnose, Downy Mildew.

### TOMATOES (5)

Early Blight, Late Blight, Anthracnose, Septorial Leaf Spot.

### SPECIFIC DIRECTIONS

- Corn** Corn Earworm
1. Dust the silk and husk necks thoroughly.
  2. Make first applications when 10 to 30% of hills show silk.
  3. Repeat at 2 to 3 day intervals for a total of 3 or more applications.
  4. Do not feed forage to dairy animals or to animals being finished for slaughter.
  5. Applications of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

**Cameraman's Note**  
Poor Copy