

Front Panel



**LILLY/MILLER**  
TOMATO-  
VEGETABLE  
DUST

ACCEPTED  
JAN 3 1983  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
this product is registered for the pesticide  
registered under  
EPA Reg. No. 600-1-74

Illustration

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl	
N-Methylcarbamate)*.....	5.0%
Rotenone.....	1.0%
Other Cube Resins.....	1.0%
Zineb (Zinc Ethylene	
Bisdithiocarbamate).....	5.0%
INERT INGREDIENTS.....	88.0%

\*SEVIN is the Reg. TM of the Union Carbide Corp.

INSECTICIDE/FUNGICIDE FOR  
TOMATOES, CARROTS, PEAS, CORN  
CUCUMBERS, CABBAGE, BEANS, MELONS

CONTAINS SEVIN\*  
ROTENONE AND ZINEB

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
See right side of panel for additional cautions

NET WEIGHT 1 LB.

Left Side Panel:

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before eating or smoking. Wash all edible portions of treated plants thoroughly before eating or processing. Do not use treated tops of vegetables for food or feed.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF ON SKIN: Wash skin thoroughly with soap and water.

IF IN EYES: Flush eyes thoroughly with water for 15 minutes. Seek medical attention immediately.

IF INHALED: Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Avoid storage of this product near feed and food products. Do not reuse empty container. Wrap container and put in trash collection.

GENERAL DIRECTIONS

Shake duster vigorously to mix air with dust thus insuring light uniform dust applications.

INSECT CONTROL

1. Dust lightly on first appearance of insect or insect damage.
2. Make complete coverage of all foliage.
3. Repeat applications every 7 to 10 days as necessary for control but be certain to observe the number of days between last application and harvest as shown in parenthesis ( ) following each crop listed.

Right Side Panel:

BEANS (7)

Cucumber Beetles, Flea Beetles, Leafhoppers, Armyworm, Stink Bugs.

BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)  
Flea Beetles, Imported Cabbageworm, Stink Bugs, Armyworm.

CARROTS (7)

Flea Beetles, Leafhoppers, Spittlebugs, Stink Bugs, Lygus Bugs.

CORN (0)

Fall Armyworm, Flea Beetles, Leafhoppers. For Corn Earworm control see Specific Directions.

CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)

Cucumber Beetles, Pickleworm, Melonworm, Flea Beetles, Leafhoppers, Squash Bugs.

HEAD LETTUCE (10), LEAF LETTUCE (14)

Flea Beetles, Leafhoppers, Armyworm, Imported Cabbageworm, Spittlebugs, Stink Bugs.

PEAS (10)

Leafhoppers, Armyworm, Grasshoppers.

TOMATOES (5), PEPPERS (1), EGGPLANT (1)

Flea Beetles, Leafhoppers, Tomato Hornworm, Tomato Fruitworm, Fall Armyworm, Stink Bugs.

DISEASE CONTROL

1. Dust lightly so as to make complete coverage of foliage at first appearance of disease or apply as a preventative before disease appears.
2. Make regular light applications at 4 to 7 day intervals until disease is under control but observe the number of days between last application and harvest as shown in parenthesis ( ) following each crop listed.

BEANS (7), PEAS (10)

Anthracoze, Downy Mildew, Rust.

BROCCOLI (7), BRUSSEL SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)  
Downy Mildew, Leaf Spot

CARROTS (7)

Early Blight, Late Blight.

Right Side Panel - Cont'd....

CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)  
Downy Mildew, Blossom Blight.

EGGPLANT (1)  
Early Blight, Late Blight, Anthracnose, Cercospora Leaf Spot.

HEAD LETTUCE (10), LEAF LETTUCE (14)  
Downy Mildew, Botrytis Slime

PEPPERS (1)  
Cercospora Leaf Spot, Anthracnose, Downy Mildew.

TOMATOES (5)  
Early Blight, Late Blight, Anthracnose, Septorial Leaf Spot.

SPECIFIC DIRECTIONS

CORN Corn Earworm

1. Dust the silk and husk necks thoroughly.
2. Make first applications when 10 to 30% of hills show silk.
3. Repeat at 2 to 3-day intervals for a total of 3 or more applications.
4. Do not feed forage to dairy animals or to animals being finished for slaughter.
5. Applications of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

---

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

---

LC-13 R

THE CHAS. H. LILLY CO.  
Portland, Oregon 97218

EPA Est 802-OR-1

EPA Reg. No. 802-474