

Colors: Yellow and white background, black type, multi-color illustration.

Type Sizes: As indicated, all other 6 pt minimum.

Front Panel:

LILLY/MILLER
Tomato-Vegetable Dust

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate)	5.0%
Rotenone	1.0%
Other Associated Resins	2.0%
INERT INGREDIENTS	92.0%

For Tomatoes, Carrots, Peas, Corn, Cucumbers, Cabbage, Beans, Melons

Contains SEVIN* brand Carbaryl Insecticide and Rotenone


*SEVIN is the registered trademark of Rhone-Poulenc for carbaryl insecticide.

Keep out of reach of children (12 pt type)

CAUTION (18 pt type)

See side panel for additional precautionary statements

NET WEIGHT 1 LB.



Side Panels:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before eating or smoking. Wash all edible portions of treated plants thoroughly before eating or processing. Do not use treated tops of vegetables for food or feed.


NOTE TO PHYSICIAN: Carbaryl Insecticide is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Induce vomiting and seek medical attention immediately. If On Skin: Wash skin thoroughly with soap and water. If In Eyes: Flush eyes thoroughly with water for 15 minutes. Seek medical attention immediately. If Inhaled: Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.



This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

(For Squeeze duster only):

With plastic cap closed, shake duster vigorously to mix air with dust. Lift plastic cap, pointing the opening towards the plants. Gently squeeze plastic tube to insure uniform, light dust application. Close cap after each use.

(For all containers):

Dust lightly on first appearance of insect or insect damage. Make complete coverage of all foliage. Repeat applications every 7 to 10 days as necessary for control but be certain to observe the number of days between last application and harvest as shown in parenthesis () following each crop listed.

BEANS (7)

Cucumber Beetles, Flea Beetles, Leafhoppers, Armyworm, Stink Bugs.

BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7)

Flea Beetles, Imported Cabbageworm, Stink Bugs, Armyworm.

CARROTS (7)

Flea Beetles, Leafhoppers, Spittlebugs, Stink Bugs, Lygus Bugs.

CORN (0)

Fall Armyworm, Flea Beetles, Leafhoppers. For Corn Earworm control see "SPECIFIC DIRECTIONS".

CUCUMBERS (5), MELONS (5), PUMPKINS (0), SQUASH (Summer and Winter) (5)

Cucumber Beetles, Pickleworm, Melonworm, Flea Beetles, Leafhoppers, Squash Bugs.


HEAD LETTUCE (10), LEAF LETTUCE (14)

Flea Beetles, Leafhoppers, Armyworm, Imported Cabbageworm, Spittlebugs, Stink Bugs.

PEAS (10)

Leafhoppers, Armyworm, Grasshoppers.

TOMATOES (5), PEPPERS (1), EGGPLANT (1)



Flea Beetles, Leafhoppers, Tomato Hornworm, Tomato Fruitworm, Fall Armyworm, Stink Bugs.

SPECIFIC DIRECTIONS

CORN (0)

Corn Earworm

1. Dust the silk and husk necks thoroughly.
2. Make first applications when 10 to 30% of hills show silk.
3. Repeat at 2 to 3 day intervals for a total of 3 or more applications.
4. Do not feed forage to dairy animals or to animals being finished for slaughter.
5. Applications of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 802-OR-1
EPA Reg. No. 802-474
90952

