

ft to
ps in
well
thers.
being
s be-
water
es in

Acre
to 20

to 20

to 20

to 15

ROOT MAGGOTS - Broccoli, Brussels Sprouts, Cauliflower,
Cabbage, Collards, Kale, Kohlrabi, Radish, Turnips
Apply as a preplanting soil application or at time
planting or transplanting.

INSECT CONTROL on HOME LAWNS and ORNAMENTAL PLANTS
use on newly seeded lawns and turfs - for the control of
Ants, Armyworms, Boxelder Bugs, Chiggers, Fleas, Grubs,
Crickets, European Earwigs, Grasshoppers, Japanese Beetles,
Grubs, Mosquitoes, White Grub, Sod Webworm, Wireworm.
*This would be at the rate of 1-1/4 lbs. per 1000 sq. ft.
Apply to cover surface uniformly in the spring or when
insects are observed to be troublesome. Immediately after treat-
ment for control of Chinch Bugs, Japanese Beetle Grubs, Sod web-
worms, Wireworms, and White Grubs, drench the lawn with water or
sprinkle heavily to make treatment more effective and to reduce
the hazard to children and pets. Do not treat when soil is wet.
Apply before rain or sprinkling.

CAUTION: Do not permit children or pets to go on treated lawn
until the insecticide has been washed off the grass into the
soil by drenching or sprinkling with water and grass has com-
pletely dried. Birds feeding on treated areas may be killed.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other
chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the
seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-4H

CHAS. H. LILLY CO.
PORTLAND, OREGON

