

Front Panel


Lawn & Turf
Weed Killer

Illustration

ACTIVE INGREDIENTS:

+ Dimethylamine 2,4-Dichloro- phenoxyacetate.....	9.41%	✓	4.46
++ Dimethylamine Salt of 2-(2-methyl-4-chloro- phenoxy) Propionic Acid.....	4.54%		4.56
+++ Dimethylamine Salt of Dicamba (3,6-dichloro-o- anisic acid).....	1.04%		1.05
INERT INGREDIENTS.....	85.01%		

KILLS
 DANDELIONS, PLANTAINS,
 JAPANESE CLOVER, CHICKWEED

with comments
 JUN 2 1965
 Rfm
 800-485


KEEP FROM FREEZING

TREATS 9600 SQ. FT. OF LAWN

Keep out of reach of children
CAUTION
 See left side panel for additional cautions

NET CONTENTS 1 QT.

BEST EXPIRES



Left Side Panel:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION. Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL CAUTIONS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Rinse thoroughly before discarding and securely wrap original container in several layers of newspaper and discard in trash.

USE PRECAUTIONS

Use separate hose-end or pressure sprayer for weed killing only.

Avoid drift of spray mist onto vegetables, grapes, flowers, ornamental plants, shrubs, shade and fruit trees, and other desirable plants. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not pour spray solutions near these plants. Do not use on Dichondra, Carpetgrass, Centipede or St. Augustine grasses. Clovers present will be damaged if sprayed.

BEST DOCUMENT AVAILABLE

Left Side Panel (Con't)

+2,4-dichlorophenoxyacetic acid equivalent 0.68 lbs per
gallon (7.82%).
Isomer Specific by AOAC method 6.D01-5.
++2-(2-methyl-4-chlorophenoxy) propionic acid equivalent
0.325 lbs. per gallon.
+++3,6-dichloro-o-anisic acid equivalent 0.075 lbs. per
gallon.
U.S. Pat. No. 3,284,186

*Please
check
equivalent
to Tuent*

Notice: Seller makes no warranty, expressed or implied,
concerning the use of this product other than indicated
on the label. Buyer assumes all risk of use and/or
handling of this material when such use and/or handling
is contrary to label instructions.

LC-5S

THE CHAS. H. LILLY CO.
Portland, Oregon 97218

EPA EST. 802-OR-1

EPA Reg. No. 802-485

BEST DOCUMENT AVAILABLE

Right Side Panel

Please read entire label. Use only as directed.
Observe all cautions and precautions.
Shake well before use

Lilly/Miller LAWN & TURF WEED KILLER controls Bedstraw, Buckhorn, Buttercup, Chickweed, Clover, Dandelion, Dock, English Daisy, Ground Ivy, Henbit, Japanese Clover (Black Medic), Knotweed, Lambsquarter, Mallow, Morningglory, Oxalis, Pigweed, Plantains, Poison Oak, Purslane, Shepherdspurse, Speedwell, Spurge, Yarrow, and most other broadleaved lawn weeds.

NEW LAWNS AND BENT GRASS TURF DIRECTIONS FOR USE

On new lawns wait until turf has been mowed two or three times and has reached a height of 2 inches before making an over-all spray treatment. Then spray with 2/3 of the dilutions given for established lawns (for example, in Hose-end Sprayers, use 5 tablespoons, fill jar to 5 gallon mark and spray 1000 sq. ft. of lawn. In pressure Sprayers, use 1 tablespoon in 1 gallon of water and spray 200 sq. ft. of lawn).

"Spot Spray" weed treatment (at half strength) may be made at any time with a small pressure sprayer. Hold the nozzle near ground level and spray the weeds only. (On Bent Grass lawns, temporary yellowing may occur.) Wait three weeks after treatment before putting new grass seed or stolons down in the bare spots.

ORNAMENTAL LAWNS & TURF (Other Than Bent Grass)

WHEN TO USE on established Lawns and Turf- spray at any time during the growing season when weeds are actively growing. Treat Chickweed in the fall or in the spring as needed. Poor weed control may result if spray is applied during drought or just before rain.

For best results, mow lawn 2 or 3 days before treating. Do not water lawn within 24 hours after treatment. Repeat application, if necessary, for hard-to-kill weeds. Observe Use Precautions.

HOW MUCH TO USE - Over-all Spray Treatment

HOSE END SPRAYERS: Use a hose-end sprayer designed for lawns. Put recommended amount of this product in jar, add water to indicated level, mix thoroughly and spray on area indicated. For example, to spray 3000 sq. ft. of lawn, measure 20 tablespoons into jar, add water to the 10 gallon level and mix thoroughly. Attach jar to hose, turn water on and spray as evenly as possible over the 3000 sq. ft.

Right Side Panel - Cont'd

Use this amount of LAWN & TURF WEED KILLER	Fill jar to (gallons)	to spray this amount of lawn or turf
2 tablespoons	1	300 sq. ft.
10 tablespoons	5	1500 sq. ft.
20 tablespoons	10	3000 sq. ft.
30 tablespoons	15	4500 sq. ft.

PRESSURE SPRAYERS - measure indicated amount of this product into tank and dilute with water according to the following chart. Use a spray pattern which gives small droplets without any fine mist and cover the area indicated.

Use this amount of LAWN & TURF WEED KILER	In this amount of water	to spray this amount of water or turf
2 tablespoons	1 gallon	300 sq. ft.
4 tablespoons	2 gallons	600 sq. ft.
6 tablespoons	3 gallons	900 sq. ft.
1 pint	16 gallons	4800 sq. ft.

NOTE: Whether using a hose-end sprayer, or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

"SPOT SPRAY" Treatment on established Lawns and Turf. To spray small area, or individual weeds, use pressure sprayer (do not use hose-end sprayer). Mix 4 table-
spoons of this product in 1 gallon of water in pressure sprayer and spray as required being certain to "wet" each weed.

BEST

Irene Boone
The Chas. H. Lilly Co.
7737 NE. Killingsworth
Portland, OR 97218

Dear Ms. Boone:

Subject: Label Revision: AOAC Method
Lilly/Miller Lawn & Turf Weed Killer
EPA Registration No. 802-485
Your submission of June 13, 1983

The amendment referred to above, submitted in connection with registration under FIFRA is acceptable, provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. Modify "Keep out of lakes streams or ponds" to read:
Do not apply directly to water
 - b. Place the ingredient referrals as a substatement immediately below the ingredient declaration.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

We are assuming that the weight of water on the Confidential Formula is a typographical error and is 2772 to be identical with previously submitted formulas.

Sincerely yours,

Richard F. Mountfort *R. F. Mountfort*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

RD:Mountfort:DCR-25858:WANG-1143C:bje:Raven:479-2013:6/23/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							