

necessary at 14 day intervals. Do not use

1 1/2 pts. per acre when insects are causing damage to dairy animals or to use of grain.

USE 2 pts. per acre on soil around plants.

ported Cabbageworm, Armyworm—USE 4 pts. per acre after seedling stage on leaf lettuce or

USE 3 pts. per acre. Apply when plants are in seedling stage in very large numbers in seed beds or spring onions.

Yellow Striped Armyworm, Cutworms—USE 4 pts. per acre. Do not feed treated vines for slaughter.

USE 3 pts. per acre. Apply to infested foliage. Do not apply within 5 days of harvest.

USE 2 pts. per acre. Apply on soil around plants. Do not apply within 5 days of harvest.

USE 2 to 3 pts. per acre. Apply on foliage around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.

USE 2 pts. per acre. Apply on soil around treated tops for food or feed.



TOXAPHENE 8E

(8 lbs. Toxaphene per gallon)

AGRICULTURAL INSECTICIDE

Active ingredients	
*Toxaphene	71.0%
Aromatic Petroleum Derivative Solvents	24.1%
Inert ingredients	4.9%
*Technical Chlorinated Camphene (Chlorine 67-69%)	

E.P.A. Reg. No. 802-486-AA

WARNING — Keep out of reach of children

May be Fatal if Swallowed! Do not breathe spray mist. May be absorbed through skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. Not for use or storage in or around the home. Do not use or store near heat or open flame. Protect from temperatures below 15° F.



— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
 PORTLAND, OREGON 97214




PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.

MIXING DIRECTIONS

1. Fill tank 1/2 to 2/3 full, add required amount of Miller's TOXAPHENE 8E, then add balance of water to fill tank.
2. Have agitator running during filling, mixing and spraying operations.
3. Do not allow mixture to stand. If mixture does not readily mix with water, but tends to separate as an oily layer, do not use as injury to plants may result.
4. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proven them to be physically compatible.

USE DIRECTIONS

1. Use the amounts per acre for specific crops in sufficient water (6 to 10 gal. by air; 20 to 50 gal. by ground equipment) to give uniform coverage of all plants.
2. Unless otherwise specified, start applications at the first sign of insects or insect injury. Repeat as needed or control, observing time limitations on specific crops.
3. Consult State Agricultural Experiment Station or State Extension Service Authorities for additional information consistent with label directions and cautions including timing and number of applications needed for your area.

USE PRECAUTIONS

Do not apply to any area not specified on the label. Avoid contamination of feed and foodstuffs. Do not apply when weather conditions favor drift from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others. This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply where run-off is likely to occur. Birds feeding on treated areas may be killed. Rinse equipment and dispose of wastes in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-6F

RECEIVED
 2/72
 THE CHAS. H. LILLY CO.
 PORTLAND, OREGON

SPECIFIC DIRECTIONS

ALFALFA—for control of Cutworms, Western Yellow Striped Armyworm, Alfalfa Weevil, Flea Beetles, Grasshoppers, Leafhoppers, Lygus Bugs—USE 2 to 3 1/2 pts. per acre.

ALFALFA SEED CROPS ONLY—for control of above insects—USE 3 1/2 pts. per acre.

GRAPE

ALFALFA—USE PRECAUTIONS

Do not graze dairy animals in treated fields. Do not graze meat animals in treated fields within 6 weeks of slaughter. Do not feed treated plants, threshings, or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not use screenings for pellets or other feed formulations.

BEANS, DRY AND QUICK FREEZE (Lima, Blackeye, Kidney, Pink)—for control of Cutworms, Darkling Ground Beetles—USE 3 pts. per acre. Apply in seedling stage.

BEANS, GREEN (Lima, Snap, String, Wax, Blackeye)—for control of Cutworms, Darkling Ground Beetles—USE 3 to 4 pts. per acre. Apply in seedling stage.

BEANS, GREEN (Lima, Snap, String, Wax, Blackeye)—for control of Bean Leafminers, Serpentine Leafminers, Flea Beetles—USE 2 pts. per acre. Make 2 applications at 15 day intervals.

BEANS—USE PRECAUTIONS

If beans are going to processor check with company representative before making pesticide applications. Do not apply to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or to animals being finished for slaughter. No time limitation on shelled beans as human food.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE/ COLLARDS, KALE, KOHL-RABI—for control of Cabbage Looper, Diamondback Moth, Imported Cabbageworm, Climbing Cutworms—USE 1 1/2 to 2 1/2 pts. per acre. Do not apply after edible parts start to form or within 30 days of harvest. On **Cabbage**, do not apply after heads start to form. On **Collards and Kale**, do not apply within 35 days of harvest.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, KOHL-RABI—for control of Cutworms (soil)—USE 2 pts. per acre. Apply on soil around plants in seedling stage. See above for Use Precautions.

CARROTS—for control of Cutworms (soil)—USE 3 pts. per acre. Apply on soil around plants in seedling stage. Do not use tops for food or feed.

CARROTS FOR SEED—for control of Lygus Bugs (adults and nymphs)—USE 2 pts. per acre. Make first application when counts reach 1/2 to 1 Lygus per seed head, usually about

25% bloom. Two to 3 applications may be necessary at 14 day intervals. Do not use tops for food or feed.

CORN, FIELD—for control of Grasshoppers—USE 1 1/2 pts. per acre when insects are causing damage, usually in seedling stage. Do not feed treated forage to dairy animals or to animals being finished for slaughter. No limitation on use of grain.

EGGPLANTS—for control of Cutworms (soil)—USE 2 pts. per acre on soil around plants as needed. Do not apply within 5 days of harvest.

LETTUCE—for control of Cabbage Looper, Imported Cabbageworm, Armyworm—USE 4 pts. per acre on foliage as needed. Do not apply after seedling stage on leaf lettuce or after heads begin to form on head lettuce.

ONIONS FOR SEED—for control of Thrips—USE 3 pts. per acre. Apply when plants are in bloom (after flowers open), and when thrips are present in very large numbers in seed head (200 per hand shake). Do not apply to green or spring onions.

PEAS—for control of Alfalfa Looper, Western Yellow Striped Armyworm, Cutworms—USE 3 pts. per acre. Do not apply after pods begin to form if pods are to be used for food, or within 7 days of harvest if pods are not to be used for food. Do not feed treated vines to dairy animals or to animals being finished for slaughter.

PEPPERS—for control of Pepper Weevil—USE 3 pts. per acre. Apply to infested foliage after weevils appear and repeat every 7 to 10 days. Do not apply within 5 days of harvest.

PEPPERS—for control of Cutworms (soil)—USE 2 pts. per acre. Apply on soil around plants as needed. Apply in late afternoon. Do not apply within 5 days of harvest.

RUTABAGAS—for control of Cabbage Looper—USE 2 to 3 pts. per acre. Apply on foliage every 7 days when insects appear. Do not use treated tops for food or feed.

RUTABAGAS—for control of Cutworms (soil)—USE 2 pts. per acre. Apply on soil around plants as needed. Apply in late afternoon. Do not use treated tops for food or feed.

TOMATOES—for control of Cutworms (soil)—USE 2 pts. per acre. Apply on soil around plants as needed. Apply in late afternoon. Do not apply within one day of harvest.

TOMATOES—for control of Tomato Hornworm, Tomato Fruitworm, Russet Mites, Flea Beetles—USE 5 pts. per acre. Do not apply within 3 days of harvest.

USE PRECAUTIONS

In California, permit for purchase or use must be obtained from County Agricultural Commissioner.

Do not graze dairy animals in treated fields. Do not graze meat animals in treated fields within 6 weeks of slaughter. Do not feed treated plants, threshings, or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter.

Apply in the early morning or late evening when pollinators are not in the fields. Ground applications are preferable. Make applications when there is little or no wind. Do not allow drift to nearby hay or forage crops.

AGC
JUN 12, 72

Insist on...

Miller's



TOXAPHENE 8E

(8 lbs. Toxaphene per gallon)

AGRICULTURAL INSECTICIDE

Active ingredients	
*Toxaphene	71.0%
Aromatic Petroleum Derivative Solvents	24.1%
Inert ingredients	4.9%
**Technical Chlorinated Camphene (Chlorine 67-69%)	

E.P.A. Reg. No. 802-486-AA

WARNING — Keep out of reach of children

May be Fatal if Swallowed! Do not breathe spray mist. May be absorbed through skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using.

Not for use or storage in or around the home.

Do not use or store near heat or open flame.

Protect from temperatures below 15° F.



Gallon

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214



PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.

MIXING DIRECTIONS

1. Fill tank 1/2 to 2/3 full, add required amount of Miller's TOXAPHENE 8E, then add balance of water to fill tank.
2. Have agitator running during filling, mixing and spraying operations.
3. Do not allow mixture to stand. If mixture does not readily mix with water, but tends to separate as an oily layer, do not use as injury to plants may result.
4. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proven them to be physically compatible.

USE DIRECTIONS

1. Use the amounts per acre for specific crops in sufficient water (6 to 10 gal. by air, 2 to 50 gal. by ground equipment) to give uniform coverage of all plants.
2. Unless otherwise specified, start applications at the first sign of insects or insect injury. Repeat as needed or control, observing time limitations on specific crops.
3. Consult State Agricultural Experiment Station or State Extension Service Authorities for additional information consistent with label directions and cautions including timing and number of applications needed for your area.

USE PRECAUTIONS

Do not apply to any area not specified on the label.

Avoid contamination of feed and foodstuffs. Do not apply when weather conditions favor drift from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply where run-off is likely to occur. Birds feeding on treated areas may be killed.

Rinse equipment and dispose of wastes in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with waste. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.