



Miller's

**4 BAIT
STATIONS INSIDE
THIS PACKAGE!**

Rat & Mouse Bait

WITH WARFARIN READY TO USE

500 GRAMS
NET WEIGHT
1.76 LBS
100% PURE



Copy of label on back of 1 lb. box ...

MILLER'S RAT AND MOUSE BAIT is especially recommended for rat and mouse infestations which are difficult to control. After initial cleanout has been accomplished, prevent reinfestation by maintaining bait stations.

DIRECTIONS FOR USE

RATS . . . Place 4 oz. to 1 lb. of bait in locations frequented by rats. Baits should be replaced as consumed. Any interruption in feeding will reduce effectiveness of control. Where continuous source of rat infestation prevails from nearby dumps or fields, permanent bait stations should be used and baits replenished as needed.

MICE . . . for mice control, place 1 to 2 tablespoons of bait at 8 to 12-foot intervals. Heap the bait in a pile as this makes a more attractive bait surface for mice. Mice are usually more difficult to control than rats with this material and satisfactory control may take a longer period of baiting.

SPECIFIC DIRECTIONS

1. When bait is placed, it is more effective if other sources of food, more attractive to rats and mice, are removed.
2. Treated baits should be placed in locations not accessible to children, pets, domestic animals and wildlife, or in tamper-proof bait boxes.
3. Assure continuous bait supply for at least 10 days for rats and 15 days for mice and thereafter until all signs of feeding have stopped. After initial cleanout has been accomplished prevent reinfestation by maintaining bait stations.
4. Remove all decomposed, mouldy or sour baits—the rats and mice will leave them alone and go to other food.
5. Bait stations give best results—they can be made as follows:

Leaning Board: Lean a board against a wall to provide a tunnel under which rats may eat. If the space beneath is kept fairly narrow, it will give the rats protection and at

the same time keep cats and other animals from reaching the bait.

Drain Tile or Sewer Pipe: These are useful around the poultry house and in similar locations. Pipe should be 4 to 6 inches in diameter and at least 24 inches long. Bait should be placed inside by means of a long-handled spoon.

CAUTION

Keep away from children, pets, domestic animals and wildlife. This material contains an anti-coagulant which if swallowed by humans, pets, domestic animals or wildlife may reduce the clotting ability of the blood and cause bleeding. In case of eating of baits, call a physician immediately. In the meantime, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear.

Note to Physician—In case of ingested bait, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Destroy empty container. Do not reuse for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON, U.S.A.