

Miller's SYSTEMIC FUNGICIDE BENOMYL provides both curative (eradicator) and protective action. The addition of a surfactant to the spray mixture enhances the curative action and improves distribution of the spray on hard-to-wet plants.

Active Ingredient:
 Benomyl (Methyl 1-[butylcarbamoyl]-2-benzimidazolecarbamate) 50%
 Inert Ingredients: 50%

CAUTION

May irritate eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water. For eyes, get medical attention.

KEEP AWAY FROM FIRE OR SPARKS.

Do not re-use container; destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-3H

Manufactured for . . .

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

Net Wt.
 1 Oz.

\$2.79



Systemic Fungicide
Benomyl

**ORNAMENTALS, LAWNS,
 STONE FRUIT, SQUASH,
 MELONS, SNAP BEANS,
 CUCUMBERS**

CAUTION: Keep out of reach of children.
 See left side panel for additional cautions
 E.P.A. Reg. No. 802-490-AA

DIRECTIONS FOR USE

ROSES, FLOWERS, ORNAMENTALS
 Powdery Mildew, Black Spot, Botrytis, Gray Mold

LAWN GRASS DISEASES

Apply on lawn grasses to the following: **Black Spot** - Repeat at 7 to 14 day intervals as long as needed. Do not graze treated areas. Do not feed clippings to livestock. **Dollar Spot** - Sclerotinia Home-Larvae - Mix 1 oz. in level tablespoons in 2 to 3 gal. water and apply evenly to 1000 sq. ft. **Large Brown Patch** - Rhizoctonia solani, **Fusarium Patch** - Fusarium nivale - Mix 1 oz. entire package in 1 to 2-1/2 gal. water and apply evenly to 500 sq. ft. When conditions are especially favorable for development of Large Brown Patch, apply spray at 5 to 7 day intervals.

**PEACHES, CHERRIES, APRICOTS, PLUMS,
 PRUNES, NECTARINES**

Brown Rot Blossom Blight - Use 1 level tablespoon in 2 gal. water. Apply as a full coverage spray at early bloom stage and repeat at full bloom. **Fruit Brown Rot** - Follow the above spray program and make 2 additional sprays beginning 3 weeks before harvest.

BEANS (Snap)

Gray Mold (Botrytis), White Rot (Sclerotinia) - Use 1 level tablespoon in 2 gal. water and apply at early bloom. Repeat application at peak bloom. Do not use within 14 days of harvest. Do not feed treated bean vines to livestock.

CUCUMBERS, MELONS, SUMMER and WINTER SQUASH
Powdery Mildew, Anthracnose, Gummy Stem Blight - Use 1 level tablespoon per 2 gal. water and apply when stems begin to run or when disease first appears. Repeat at 7 to 10 day intervals as needed to control disease.

DIRECTIONS FOR USE

IMPORTANT: Occasionally shake or agitate sprayer to insure even mixture of spray.

ROSES, FLOWERS, ORNAMENTALS
 Powdery Mildew, Black Spot, Botrytis, Gray Mold: Use 1 level tablespoon

Miller's SYSTEMIC FUNGICIDE BENOMYL provides both curative (eradicator) and protective action. The addition of a surfactant to the spray mixture enhances the curative action and improves distribution of the spray on hard-to-wet plants.

Net Wt.
 3 Oz.

\$5.49