

Miller's Systemic Fungicide BENOMYL provides a protective and preventive action. It controls a wide range of fungal diseases on ornamentals, lawns, and fruit trees. It is also effective against many of the most common lawn diseases.

**Application:** Benomyl is applied as a spray or dust to the foliage of plants.

**Directions:** See the label for directions on the use of Benomyl on various plants.

**CAUTION**

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

**KEEP AWAY FROM FIRE OR SPARKS**

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

**THE CHAS. H. LILLY CO.**  
PORTLAND, OREGON

Net Wt.  
1 Oz.

\$2.79

**Miller's**  
**Systemic Fungicide**  
**Benomyl**

**ORNAMENTALS, LAWNS,  
STONE FRUIT, SQUASH,  
MELONS, SNAP BEANS  
CUCUMBERS**

See left side panel for additional cautions  
EPA Reg. No. 60249C-AA

**DIRECTIONS FOR USE**

**ROSES, FLOWERS, ORNAMENTALS**  
Powdery Mildew, Black Spot, Botrytis Gray Mold - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**LAWN GRASS DISEASES**  
Dollar Spot, Brown Patch, Large Brown Patch, Fusarium Patch - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**PEACHES, CHERRIES, APRICOTS, PLUMS, PRUNES, NECTARINES**  
Brown Rot Blossom Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**BEANS (Snap)**  
Gray Mold, Botrytis, White Rot, Scab of Bean - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**CUCUMBERS, MELONS, SUMMER and WINTER SQUASH**  
Powdery Mildew, Anthracnose, Gummy Stem Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

Net Wt.  
3 Oz.

\$5.49

**Miller's**  
**Systemic Fungicide**  
**Benomyl**

**ORNAMENTALS, LAWNS, STONE  
FRUIT, APPLES, PEARS, STRAW-  
BERRIES, CUCUMBERS, SQUASH,  
MELONS, SNAP BEANS,  
BULBS**

**CAUTION!** Keep out of reach of children.  
See left side panel for additional cautions.  
EPA Reg. No. 60249C-AA

**DIRECTIONS FOR USE**

**IMPORTANT!** Occasionally shake or agitate spray container prior to use of spray.

**ROSES, FLOWERS, ORNAMENTALS**  
Powdery Mildew, Black Spot, Botrytis Gray Mold - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**LAWN GRASS DISEASES**  
Dollar Spot, Brown Patch, Large Brown Patch, Fusarium Patch - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**PEACHES, CHERRIES, APRICOTS, PLUMS, PRUNES, NECTARINES**  
Brown Rot Blossom Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**BEANS (Snap)**  
Gray Mold, Botrytis, White Rot, Scab of Bean - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**CUCUMBERS, MELONS, SUMMER and WINTER SQUASH**  
Powdery Mildew, Anthracnose, Gummy Stem Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**BULBS**  
Fusarium and Penicillium Rots on Easter Lily, Tulip, Gladiolus, Daffodil, Iris - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**STRAWBERRIES**  
Gray Mold, Powdery Mildew, Leaf Scorch, Botrytis, Gummy Stem Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**APPLES-PEARS**  
Powdery Mildew, Scab, Botrytis, Fire Blight, Gummy Stem Blight - Use 1 level tablespoon (2.8 level teaspoons) per gallon of water. Apply at 7 to 14 day intervals when disease first appears and repeat at 7 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

Miller's Systemic Fungicide BENOMYL provides a protective and preventive action. It controls a wide range of fungal diseases on ornamentals, lawns, and fruit trees. It is also effective against many of the most common lawn diseases.

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

**CAUTION**

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

**KEEP AWAY FROM FIRE OR SPARKS.**

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

Miller's Systemic Fungicide BENOMYL is a systemic fungicide. It is highly effective against a wide range of fungal diseases. It is also effective against many of the most common lawn diseases.

**THE CHAS. H. LILLY CO.**  
PORTLAND, OREGON

ACCEPTED

ACCEPTED

OCT 25 1969

UNDER THE FEDERAL INSTITUTE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO. 8-27-69

# SA BRAND 50<sup>ai</sup> TOMATO DUST

### ACTIVE INGREDIENTS:

Copper (from basic copper sulfate) . . . . .	7.00%
INERT INGREDIENTS . . . . .	93.00%
	100.00%

### A FUNGICIDE DUST FOR USE ON

**TOMATOES and POTATOES** for the control of early and late blights as an aid in controlling bacterial spot. Dust plants thoroughly at weekly intervals and following rains. For potatoes make 30 to 60 lbs. per acre applications beginning when the plants are 6 inches high. For tomatoes make 30 to 60 lbs. per acre applications beginning when plants are set in the field.

**CUCUMBERS, SQUASH, WATERMELONS and CANTALOUPEs** for the control of downy mildew. Dust thoroughly, but lightly, using 15 to 40 lbs. per acre. Apply at 10-day intervals and following rains. Heavy applications, applications to tender foliage, and applications to wet foliage can cause injury to cucurbits.

**PEPPER** as an aid in the control of bacterial spot, apply at weekly intervals and if possible preceding heavy rains, using 20 to 50 lbs. per acre applications.

There is no time limitation on the use of this dust. Remove excess residue at harvest. Consult proper State Extension specialist to determine if local recommendations are consistent with this label before using. Do not exceed indicated dosages. May cause injury to tender plant parts particularly when applications are excessive, uneven, or when foliage is wet. Control of the indicated diseases may not be satisfactory during periods of frequent or heavy rainfall, when disease infestations are extremely severe, or when applications are uneven, inadequate, or poorly timed.

Do not apply or allow drift to corn, grain sorghum, or other copper sensitive plants. Do not apply when conditions favor drift from treated area.

Destroy empty container.

### Keep Out of Reach of Children

**CAUTION** Harmful if swallowed. Avoid contact with skin, eyes, and clothing. Do not contaminate feed or foodstuffs.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions given herewith.

## Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.  
E. S. D. A. Reg. No. 8296 B

NET WT. 5 LBS.

Please note: Inert ingredient statement, packing sizes, and label size on other side.