


DANGER - POISON - PRECAUTIONS ☠


STOP - READ THE LABEL

DON'T SWALLOW




POISONOUS IF SWALLOWED
Even in small amounts!

DON'T TOUCH



POISONOUS BY SKIN CONTACT
Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothes (liquid goes through clothes).

DON'T BREATHE



POISONOUS IF BREATHED
Poisonous if vapors or mists from sprays are breathed. Vapors are not visible. Never work with methyl parathion or in methyl parathion treated areas without protective clothing and equipment.

POISONOUS TO FISH AND WILDLIFE: Toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where run-off is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

POISON

See Side Panels
for Antidote &
Precautions



DANGER

Keep Out of
Reach of
Children

EMULSIFIABLE
LIQUID

ACCEPTED

Miller's

METHYL PARATHION 4E

(4 lbs. Methyl Parathion per gallon)

ORGANOPHOSPHORUS INSECTICIDE

**AGRICULTURAL CHEMICAL
NOT FOR HOME USE**

Active ingredients:

O,O-dimethyl O-p-nitrophenyl
phosphorothioate* 45.50%
Aromatic Petroleum Derivative Solvents . 47.62%
Inert ingredients: 6.88%

*Methyl Parathion
COMBUSTIBLE—Keep away from heat &
open flame.

Read Label Folder for additional use pre-
cautions, directions for use, recommenda-
tions and container disposal.

E.P.A. Reg. No. 802-498-AA

See side panels for poison precautions,
symptoms, first aid treatment, information
for physician and posting treated areas.

NOTICE: Follow directions carefully. Timing and
method of application, weather and crop condi-
tions, mixtures with other chemicals not specifi-
cally recommended, and other influencing factors
in the use of this product are beyond the control
of seller. Buyer assumes all risks of use, storage
or handling of this material not in strict accord-
ance with directions given herewith.

NET CONTENTS: Gallon(s)

Manufactured for . . .
— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON

Miller's

SEVIN* 5B

(5% SEVIN* BAIT)

AGRICULTURAL INSECTICIDE

Aug 29, 72

NET WT. 50 LBS.

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214

Active ingredient

Carbaryl (1-naphthyl N-methylcarbamate)* 5%

Inert ingredients 95%

*SEVIN is the Reg. T.M. of the Union Carbide Corp. for
1-naphthyl N-methylcarbamate. U.S.Pat. No. 2,803,478.

E.P.A. Reg. No. 802-493-AA

CAUTION — Keep out of reach of children

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid contact with skin. Wash hands before eating or smoking. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

USE PRECAUTIONS

Avoid contamination of foodstuffs, feed, feeding troughs and watering receptacles. Keep out of reach of animals.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply when honey bees are foraging in the field.

Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

ARMYWORMS, CRICKETS, CUTWORMS, DARKLING GROUND BEETLES, GRASSHOPPERS and SOWBUGS

For control of the above insects, use the pounds per acre given below for the specific crops. Broadcast applications may be made with either ground equipment or by aircraft. Repeat treatments as necessary, except where otherwise noted.

† Do not apply closer to harvest than the number of days indicated for the specific crop.

CROP	†Days	Pounds per acre
<u>Asparagus, Strawberries</u>	1	40
On Asparagus, do not repeat application within 3 days.		
<u>Beet Greens, Chinese Cabbage, Collards, Endive (Escarole), Kale, Leaf Lettuce, Parsley, Spinach, Sugar Beets, Swiss Chard, Turnip Greens</u>	14	40
<u>Beet (roots), Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Horseradish, Head Lettuce, Parsnips, Radishes, Rutabagas, Turnip (roots)</u> ..	3	40
<u>Beans, Carrots, Corn for Forage, Eggplants, Okra, Peas, Peppers, Tomatoes</u>	0	40
<u>Alfalfa, Clover, Peanuts</u>	0	30
<u>Cucumbers, Melons, Squash (Summer & winter)</u>	0	20
<u>Cotton</u>	0	40

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors to the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.