PM 19 802-493 JUN - 2 1994 Under the Federal Insecticide. Fungicide, and Rodenticide Act. as amonded, for the pesticide LILLY/MILLER registered under EPA Reg. No. 802 - 493 GRASSHOPPER & EARWIG BAIT For Use In Vegetable & Flower Gardens. Also Kills Ants, Cutworms, Crickets & Others. ACTIVE INGREDIENT: Carbaryl (1-naphthyl N-methylcarbamate) 5% 95% Keep out of reach of children CAUTION See side panel for additional precautionary statements. NET WEIGHT 2 LBS. PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: May be harmful' if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. ENVIRONMENTAL HAZARDS This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. GRASSHOPPER, EARWIG & SOWBUG BAIT may be broadcast in vegetable gardens, on lawns, and in ornamental plantings. It can protect plants from grasshoppers, earwigs, ants, cutworms, armyworms, crickets, darkling ground beetles, centipedes, millipedes and sowbugs. DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household use only. GENERAL DIRECTIONS 1. Broadcast bait evenly on first appearance of insect or insect damage. See Specific Directions given below for vegetable gardens, flowers, shrubs and lawns. 2. Repeat applications as necessary for control, but for vegetables observe the number of days before harvest for last application noted by each crop (for instance on Cabbage, Broccoli, Brussels Sprouts, and Cauliflower, you may apply this bait up to 3 days before picking or

03

harvest). 3. Repeat applications if rain occurs within 24 hours after application of bait. One cup of GRASSHOPPER & EARWIG BAIT equals approximately 3 4. ounces. SPECIFIC DIRECTIONS Vegetable Gardens: Use GRASSHOPPER & EARWIG BAIT to Control Armyworms, Crickets, Cutworms, Darkling Ground Beetles, Grasshoppers, Sowbugs, Earwigs, Ants, Centipedes, and Millipedes. Make Broadcast applications on soil to control these insects on listed vegetables. Repeat applications as necessary. Do not apply closer to harvest than the number of days shown in parenthesis for the specific vegetable crop. Oz. per ۲ 250 sq. ft. Crop Asparagus, Strawberries (1) 4 oz. On Asparagus, do not repeat application within 3 days Beet Greens, Chinese Cabbage, Collards, Endive (Escarole), Kale, Leaf Lettuce, Parsley, Spinach, Sugar Beets, Swiss Chard, Turnip Greens (14) . . . 4 oz. Beets (roots), Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Horseradish, Head Lettuce, Parsnips, Radishes, Rutabagas, Turnip (roots)(3) 4 oz. Beans, Carrots, Corn, Eggplants, Peas, Peppers, Tomatoes (0) 4 oz. Cucumbers, Melons, Squash (Sum-_ mer & Winter)(0) 2 oz. LAWNS, FLOWERS, SHRUBS AND OTHER ORNAMENTALS

To control armyworms, crickets, cutworms, darkling ground beetles, grasshoppers, sowbugs, earwigs, ants, centipedes and millipedes.

Cutworms and earwigs are active at night and spend the daytime just below the soil surface, in cracks and crevices or underneath leaf litter and rocks. Sowbugs prefer damp areas.

For protection of ornamental plants: Apply GRASSHOPPER & EARWIG BAIT

around ornamentals in shady, protected areas, at the base of walls and fences, and in 2 to 4 inches wide bands along the edge of walkways and patios. Use 2 to 4 oz. of GRASSHOPPER & EARWIG BAIT per 250 sg. ft. For best results be sure that area to be treated is damp.

For use on lawns: To control armyworms, crickets, cutworms, darkling ground beetles, grasshoppers, sowbugs, earwigs, ants, centipedes and millipedes. Mow lawn, remove clippings and thoroughly water. Apply GRASSHOPPER & EARWIG BAIT at the rate of 4 to 8 oz. per 500 sq. ft.

STORAGE AND DISPOSAL

ι

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Seller makes no'warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218 EPA Reg. No. 802-493 EPA Est. 802-WA-1 46666