


LILLY/MILLER  
GRASSHOPPER & EARWIG BAIT

For Use In Vegetable & Flower Gardens. Also Kills Ants, Cutworms, Crickets & Others.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) . . . . 5%

INERT INGREDIENTS . . . . . 95%



Keep out of reach of children

CAUTION

See side panel for additional precautionary statements.

NET WEIGHT 2 LBS. (908 g)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

GRASSHOPPER & EARWIG BAIT may be broadcast in vegetable gardens, on lawns, and in ornamental plantings. It can protect plants from grasshoppers, earwigs, ants, cutworms, armyworms, crickets, darkling ground beetles, centipedes, millipedes and sowbugs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household use only.

One cup of GRASSHOPPER & EARWIG BAIT equals approximately 3 ounces.

VEGETABLES AND FRUIT- To kill Armyworms, Crickets, Cutworms, Darkling Ground Beetles, Grasshoppers, Sowbugs, Earwigs, Ants, Centipedes and Millipedes. Broadcast lightly on soil surface around plants at the rate indicated for each crop. Do not allow bait to contact foliage or edible part of crops. Observe the number of days before harvest for last application as noted in parentheses following each crop.

Asparagus (1): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days when applications are to spears; apply as often as every 3 days when applications are to ferns. Make no more than 3 applications prior to harvest and 2 additional applications to ferns only.

Snap Beans (fresh), Peas (fresh) (all 3): Use 4 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 4 applications per year.

Beets (roots), Carrots, Radish, Parsnips, Rutabaga, Turnips (roots) (7): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi (3): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Chinese Cabbage, Kale, Collards (14): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Sweet Corn (2): Use 5 tsp per 25 sq. ft. Reapply as often as every 3 days. Make no more than 8 applications per year.

Cucumber, Melon, Squash (3): Use 3 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 6 applications per year.

Lettuce (head and leaf), Endive, Celery, Dandelion, Parsley, Swiss Chard, Hanover Salad, Rhubarb, Beets (Tops), Turnip (Tops) (15): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Potatoes (7): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Strawberries (7): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 5 applications per year.

Tomatoes, Peppers, Eggplant (3): Use 5 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 4 applications per year.

### LAWNS, FLOWERS, SHRUBS AND OTHER ORNAMENTALS

To control armyworms, crickets, cutworms, darkling ground beetles, grasshoppers, sowbugs, earwigs, ants, centipedes and millipedes. Cutworms and earwigs are active at night and spend the daytime just below the soil surface, in cracks and crevices or underneath leaf litter and rocks. Sowbugs prefer damp areas.

For protection of ornamental plants: Apply GRASSHOPPER & EARWIG BAIT around ornamentals in shady, protected areas, at the base of walls and fences, and in 2 to 4 inch wide bands along the edge of walkways and patios. Use 2 to 4 oz. of GRASSHOPPER & EARWIG BAIT per 250 sq. ft. For best results be sure that area to be treated is damp.

For use on lawns: To control armyworms, crickets, cutworms, darkling ground beetles, grasshoppers, sowbugs, earwigs, ants, centipedes and millipedes. Mow lawn, remove clippings and thoroughly water. Apply GRASSHOPPER & EARWIG BAIT at the rate of 4 to 8 oz. per 500 sq. ft.

343

**STORAGE AND DISPOSAL**

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

THE GARDEN GROW COMPANY  
PO BOX 100  
INDEPENDENCE, OREGON 97351  
EPA Reg. No. 802-493  
EPA Est. 35497-OR-1