



25 Lbs.

Contains 5% Malathion Insecticide

FASCO POULTRY DUST M is an insecticidal and miticidal dust containing 5°, Malathion for use against certain insect and mite pests of poultry.

## DIRECTIONS

Use contrary to these directions may cause excessive residues, unsatisfactory control or other unintended consequences.

POULTRY (Direct Application), Chickens, Ducks, Geese, Turkeys—For the control of Northern fowl mites, poultry lice, and as a supplement to premise treatments for chicken red mites, dust individual birds. "Use I lb. per 125 birds. Do not exceed this rate". Apply dust by means of a shaker can or with a rotary hand duster. Repeat as necessary. Avoid contamination of food and water containers.

POULTRY HOUSES: For the control of Northern fowl mites, chicken red mites, and poultry lice, apply I lb. per 40 square feet of litter and floor space. Ripeat as necessary. Avoid contaminating feed, feeding troughs, and water fountains with dust.

ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate) INERT INGREDIENTS

Trade Marks Req. U.S. Pitent Office USDA Reg No. 2312 665

CAUTION: Keep out of the reach of children. See side panel for additional cautions.

FGPA 100

BY WT. 5.0° 。 95.0° ° Total — 100.0°

breathing of dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

CAUTION

May be harmful if swallowed. Avoid prolonged

## CONDITIONS OF SALE

Al statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.