

SYSTEMIC FUNGICIDE BENOMYL provides (systemic and protective) and protective action. The wetting agent in the spray mixture enhances penetration and improves distribution of the spray on hard-to-wet plants.

Active Ingredient:  
 Benomyl (Methyl 1-[butylcarbamoyl]-2-benzimidazolecarbamate) . . . . . 50%  
 Inert Ingredients: . . . . . 50%

**CAUTION**

May irritate eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water. For eyes, get medical attention.

**KEEP AWAY FROM FIRE OR SPARKS.**

Do not re-use container; destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-3H

Manufactured for . . .

**THE CHAS. H. LILLY CO.**  
 PORTLAND, OREGON

Net Wt.  
 3 Oz.

\$5.49

**Systemic Fungicide**  
**Benomyl**

**DIRECTIONS FOR USE**

**IMPORTANT:** Occasionally shake or agitate sprayer to insure even mixture of spray.

**ROSES, FLOWERS, ORNAMENTALS**  
 Powdery Mildew, Black Spot, Botrytis Gray Mold: Use 1 level tablespoon in 2 gal. water (1-1/2 level teaspoons per gal.). Mix thoroughly. Begin spray applications when disease first appears and repeat at 10 to 14 day intervals during growing season. Shorten intervals between applications during humid or rainy weather.

**LAWN GRASS DISEASES**  
 Spray on lawn areas as directed below. Begin applications at first appearance of disease and repeat at 10 to 14 day intervals as long as needed. Do not graze treated areas. Do not feed clippings to livestock. Dollar Spot (Sclerotinia Homecarpa) - Mix 1 oz. (6 level tablespoons) in 2 to 5 gal. water and apply evenly to 1000 sq. ft. Large Brown Patch (Rhizoctonia solani), Fusarium Patch (Fusarium nivale) - Mix 2 oz. (12 level tablespoons) in 2 to 5 gal. water and apply evenly to 1000 sq. ft. When conditions are especially favorable for development of Large Brown Patch, apply spray at 5 to 7 day intervals.

**PEACHES, CHERRIES, APRICOTS, PLUMS, PRUNES, NECTARINES**  
 Brown Rot Blossom Blight - Use 1 level tablespoon in 2 gal. water. Apply as a full coverage spray at early bloom stage and repeat at full bloom. Fruit Brown Rot - Follow the above spray program and make 1 to 2 additional sprays beginning 3 weeks before harvest.

**BEANS (Snap)**  
 Gray Mold (Botrytis), White Rot (Sclerotinia) - Use 1 level tablespoon in 2 gal. water and apply at early bloom. Repeat application at peak bloom. Do not use within 14 days of harvest. Do not feed treated bean vines to livestock.

**CUCUMBERS, MELONS, SUMMER and WINTER SQUASH**  
 Powdery Mildew, Anthracnose, Gummy Stem Blight - Use 1 level tablespoon per 2 gal. water and apply when stems begin to run or when disease first appears. Repeat at 7 to 10 day intervals as needed to control disease.

**BULBS**  
 Fusarium and Penicillium Rots on Easter Lilly, Tulp, Gladiolus, Daffodil, Iris - Use 2 level tablespoons per gal. water. Soak cleaned bulbs for 15 - 30 min. in warm (80°-85°F) dip, preferably within 48 hours after digging. Dry bulbs after treatment.

**STRAWBERRIES**  
 Gray Mold, Powdery Mildew, Leaf Scorch, Leaf Blight, Leaf Spot - Use 1 level tablespoon in 2 gal. water. Apply to give thorough coverage at beginning of bloom and repeat at full bloom. Additional applications at 10 - 14 day intervals may be made if needed.

**APPLES-PEARS**  
 Powdery Mildew, Scab - Use 1 level teaspoon in 1 gal. water. Make first application just before blooming and repeat at 7 to 14 day intervals as needed for control. Note: Miller's Benomyl may cause russeting on Golden Delicious, Rome and Stayman varieties of apples.

**ACCEPTED**  
 CAUTION: Keep out of reach of children.  
 See left side panel for additional cautions.  
 E.P.A. Reg. No. 002-490-AA  
 4/10/73

002-490