

DANGER PRECAUTIONS IN USING

Vapona Insecticide concentrate and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings.) Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion be prepared from the concentrate that will be used immediately. Mix the concentrate into water or oil outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away. Do not contaminate feed, water or foodstuffs. Do not apply when weather conditions favor drift of spray from areas treated. Rapidly absorbed through skin. Replace gloves frequently and destroy used gloves. Wash thoroughly with soap and water after handling.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by breaking or by punching holes in them and burying with wastes.

In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinsing with water.

DIRECTIONS FOR USE

MUSHROOM HOUSES - Phoride Flies (Manure Fly)

Use 5 oz. in 12 1/2 gallons of water to make a 0.5 solution.

Use 1 pint of the 0.5 solution per 100 square feet.

Apply with a brush or as a coarse spray, to inside walls and doors, ventilators and cracks in buildings before new mushrooming into production.

NOTICE: This product is highly toxic to fish and wildlife. Do not use in or near streams, rivers, lakes, ponds, or other bodies of water. Do not use in or near areas where fish and wildlife are present. Do not use in or near areas where crops are raised. Do not use in or near areas where children or animals are present. Do not use in or near areas where food is prepared or stored. Do not use in or near areas where food is consumed. Do not use in or near areas where food is sold. Do not use in or near areas where food is distributed. Do not use in or near areas where food is transported. Do not use in or near areas where food is stored. Do not use in or near areas where food is consumed. Do not use in or near areas where food is sold. Do not use in or near areas where food is distributed. Do not use in or near areas where food is transported.



VAPONA®

INSECTICIDE

DANGER--POISON

Keep out of reach of children.

See right side panel for additional warnings and antidote

Active ingredients	
2,2-Dichlorovinyl Dimethyl Phosphate	9
Insecticidally active related compounds	1
Total Active Ingredients	10

EPA Reg. No. 802-497 AA

VAPONA is the registered trademark of The Chas. H. Lilly Co.

Net Contents Five Gallons

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214



WARNING

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES.

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.
Do not get in eyes, on skin or on clothing. Do not breathe fumes. Wear natural rubber gloves, waterproof protective clothing and goggles when handling the concentrate to prepare emulsions. During application indoors wear a full face mask of a type passed by the U.S. Bureau of Mines for DDVP, VAPONA Insecticide protection.

Keep all unprotected persons out of operational area.
In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.



ANTIDOTE AND FIRST AID TREATMENT

Atropine is the emergency antidote for Vapona Insecticide poisoning. It is available in an ampule containing an adequate supply of 1.0% aqueous solution for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat, or by giving warm salt water, 1 tablespoon of salt to a glass of water. Rinse with warm salt water until vomit fluid is clear. Have patient lie down and keep warm. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 15 minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1.0% aqueous atropine tablets. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. See below.

Keep patient quiet and quiet. If patient is not breathing, administer artificial respiration immediately. If patient is not breathing, transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, difficulty in breathing, blurred vision, dry mouth, dry skin, ataxia, tachycardia, and vomiting. Administer atropine immediately.

TREATMENT: Atropine is the antidote. Therapeutic antagonism is not indicated. Supportive therapy with oxygen, if there are respiratory depression, and sodium bicarbonate should be administered if the patient is acidotic. Do not use emetics. Do not use cathartics. Do not use diuretics. Do not use vasodilators. Do not use sedatives. Do not use hypnotics. Do not use tranquilizers. Do not use anticholinergics. Do not use antispasmodics. Do not use anticonvulsants. Do not use antibiotics. Do not use antifungals. Do not use antivirals. Do not use antiparasitics. Do not use antineoplastic agents. Do not use immunosuppressants. Do not use cytotoxic agents. Do not use cytostatics. Do not use cytoreductives. Do not use cytotoxic antibiotics. Do not use cytotoxic antineoplastic agents. Do not use cytotoxic immunosuppressants. Do not use cytotoxic antineoplastic agents. Do not use cytotoxic immunosuppressants. Do not use cytotoxic antineoplastic agents. Do not use cytotoxic immunosuppressants.

PREVENTION: Do not use in or near areas where children or animals are present. Do not use in or near areas where food is prepared or stored. Do not use in or near areas where food is consumed. Do not use in or near areas where food is sold. Do not use in or near areas where food is distributed. Do not use in or near areas where food is transported. Do not use in or near areas where food is stored. Do not use in or near areas where food is consumed. Do not use in or near areas where food is sold. Do not use in or near areas where food is distributed. Do not use in or near areas where food is transported.