



SUPER RICH CRABGRASS KILLER & LAWN FOOD

Lilly's SUPER RICH CRABGRASS KILLER & LAWN FOOD is a complete lawn fertilizer in addition to a pre-emergence control of Crabgrass and Annual Bluegrass (*Poa annua*). To be effective as a pre-emergence weed control, this product must be evenly applied prior to the germination of the weed seeds, as the mature plants are not affected. Do not plant lawn seed in treated area until at least 60 days after treatment.

DIRECTIONS

A. FOR CRABGRASS CONTROL:

1. Apply during April, May or June before crabgrass plants emerge.
2. Use 20 lb. per 2,500 sq. ft. by selecting the setting for your spreader from row A in the chart below.
3. Water area after applying material.

B. FOR ANNUAL BLUEGRASS CONTROL:

1. Apply during April or May AND AGAIN in September or October.
2. Use 20 lb. per 2,000 sq. ft. by selecting the setting for your spreader from row B in the chart below.
3. Water area after applying material.

SPREADER SETTINGS

APPLICATION RATE	LILLY SPREADER Model 1002	LAWN BEAUTY Models 10 & 24	SCOTT'S Models 53 & 75
A CRABGRASS CONTROL	8	6½	8½
B ANNUAL BLUEGRASS	8½	7	9

NOTICE:

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial setting, walking speed, wind, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Lilly Chemical Co. nor the retail dealer can guarantee 100% coverage as indicated by the recommended 100% settings.

CAUTION

Harmful if swallowed. Avoid contact with eyes and skin. In case of contact, flush eyes with plenty of water. Wash off with plenty of water.

Do not store near food and feed stuffs, cereals, or other pesticides, in order to prevent contamination.

Do not plant lawn seed in treated area until at least 60 days after treatment. Do not apply to newly seeded lawns (with C) seeds). Do not use on Blueberries. Application should not be made to Colchester or Toronto plants.

Store in a dry place.

Do not graze livestock on treated areas or feed plant refuse or grass clippings to livestock.

Do not reuse empty container. Destroy it by burying with waste. Dispose of wastes by burying in non-crop lands away from water supplies.

CRABGRASS KILLER & LAWN FOOD



SUPER RICH CRABGRASS KILLER & LAWN FOOD

Lilly's SUPER RICH CRABGRASS KILLER & LAWN FOOD is a complete lawn fertilizer in addition to a pre-emergence control of Crabgrass and Annual Bluegrass (*Poa annua*). To be effective as a pre-emergence weed control, this product must be evenly applied prior to the germination of the weed seeds, as the mature plants are not affected. Do not plant lawn seed in treated area until at least 60 days after treatment.

DIRECTIONS

A. FOR CRABGRASS CONTROL:

1. Apply during April, May or June before crabgrass plants emerge.
2. Use 20 lb. per 2,500 sq. ft. by selecting the setting for your spreader from row A in the chart below.
3. Water area after applying material.

B. FOR ANNUAL BLUEGRASS CONTROL:

1. Apply during April or May AND AGAIN in September or October.
2. Use 20 lb. per 2,000 sq. ft. by selecting the setting for your spreader from row B in the chart below.
3. Water area after applying material.

SPREADER SETTINGS

APPLICATION RATE	LILLY SPREADER Model 1002	LAWN BEAUTY Models 18 & 24	SCOTT'S Models 55 & 75
A CRABGRASS CONTROL	8	6½	8½
B ANNUAL BLUEGRASS	8½	7	9

NOTICE:

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chem. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

CAUTION

Harmful if swallowed. Avoid contact with eyes and skin. In case of contact, flush eyes with plenty of water. Wash after handling or using.

Do not store near feed and feedstuffs, seeds, or other pesticides, in order to prevent contamination.

Do not plant lawn seed in treated area until at least 60 days after treatment. Do not apply to newly seeded lawns (wait 60 days). Do not use on Dichondra. Application should not be made to Coltansey or Toronto Plants.

Store in a dry place.

Do not graze livestock on treated areas or feed plant refuse or grass clippings to livestock.

Do not reuse empty container. Destroy it by burying with waste. Dispose of wastes by burying in non-crop lands away from water supplies.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen N (Minimum) 15.0% Soluble Potash (K₂O) (Minimum) ... 6.0%

grass and Annual Bluegrass (Poa annua). To be effective as a pre-emergence weed control, this product must be evenly applied prior to the germination of the weed seeds, as the mature plants are not affected. Do not plant lawn seed in treated area until at least 60 days after treatment.

DIRECTIONS

A. FOR CRABGRASS CONTROL:

1. Apply during April, May or June before crabgrass plants emerge.
2. Use 20 lb. per 2,500 sq. ft. by selecting the setting for your spreader from row A in the chart below.
3. Water area after applying material.

B. FOR ANNUAL BLUEGRASS CONTROL:

1. Apply during April or May AND AGAIN in September or October.
2. Use 20 lb. per 2,000 sq. ft. by selecting the setting for your spreader from row B in the chart below.
3. Water area after applying material.

SPREADER SETTINGS

APPLICATION RATE	LILLY SPREADER Model 1682	LAWN BEAUTY Models 18 & 24	SCOTT'S Models 53 & 75
A CRABGRASS CONTROL	8	6½	8½
B ANNUAL BLUEGRASS	8½	7	9

NOTICE:

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

CAUTION

Harmful if swallowed. Avoid contact with eyes and skin. In case of contact, flush eyes with plenty of water. Wash after handling or using.

Do not store near feed and feedstuffs, seeds, or other pesticides, in order to prevent contamination.

Do not plant lawn seed in treated area until at least 60 days after treatment. Do not apply to newly seeded lawns (wait 60 days). Do not use on Dichondra. Application should not be made to Colansey or Toronto Bams.

Store in a dry place.

Do not graze livestock on treated areas or feed plant refuse or grass clippings to livestock.

Do not reuse empty container. Destroy it by burying with waste. Dispose of wastes by burying in non-crop lands away from water supplies.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen N (Minimum) 15.0%	Soluble Potash (K ₂ O) (Minimum) 6.0%
8% (Minimum) water soluble	(Derived from Muricite of Potash)
7% (Minimum) water insoluble	Calcium as S (Minimum) 5.0%
(Derived from Ureaform, lignite	(Derived from Ammonium
Leather Meal, Urea and	Sulphate and Sulfa Sol)
Ammonium Sulphate)	Iron as Fe (Minimum) 1.5%
Available Phosphoric Acid (P ₂ O ₅)	(Derived from Sulfa Sol)
(Minimum) 4.0%	
(Derived from Superphosphate)	

WARRANTY:

The use of this material being subject to conditions beyond our knowledge and control, the seller makes no warranty of any kind, express or implied, concerning the effects of this product, when not used in accordance with directions.

NET WEIGHT 20 LBS.

Manufactured by **THE CHAS. H. LILLY CO.**, Portland, Oregon 97214