

Copy of label on Printed bag ...  
Front ...

12/20/72

36



# SUPER RICH FEED & WEED

## 12-2-3

ACCEPTED

Comments in letter

JAN 19 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 2327-10 SUBJECT  
TO ATTACHED COMMENTS

changed to: 802-505

### HERBICIDE GUARANTEE

#### ACTIVE INGREDIENTS:

- 2,4-Dichlorophenoxyacetic acid ..... 0.25%
- 2-(2-methyl-4-chlorophenoxy) propionic acid ..... 0.15%
- Dimethylamine salt of Dicamba (2-methoxy-3,6-dichlorobenzic acid) .. 0.09%

INERT INGREDIENTS ..... 99.51%

Contains TRIMEC™ U.S. Patent No. 3,284,168 T.M.P.B.I. K.C.K.  
DANVEL-D is the registered trademark of the Velocel Chemical Co.

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**SEE BACK PANEL FOR ADDITIONAL CAUTION**

E.P.A. Reg. No. 2327-10-AB-602

## NET WEIGHT 20 LBS.

Back Panel of label on Printed Bag ...

Lilly's Feed & Weed is a real work saver. Through its XF (Extended Feeding) ingredients, it feeds the lawn for weeks, thus reducing the frequency of fertilizing and mowing. TRIMEC, Lilly's effective weed killer, saves the hard work of digging and pulling weeds. Lilly's exclusive PENATURF relieves the surface tension of compact or dense soils to insure that water and food penetrates deep down to the feeding grass roots. This means less watering.

**SUPER RICH FEED & WEED** effectively controls a broad range of lawn weeds, including dandelion, plantain, chickweed, (common & mouse ear), knotweed, clover, Japanese clover, henbit, buttercup, spurge, puncture vine, lawn daisy and other similar weeds.

Apply LILLY'S FEED & WEED any time weeds are growing actively—spring, summer or fall. Applications to moist foliage and soil when temperature is between 50° and 80° F. are the most effective. Applications during hot weather (over 80° F.) or on dry soil are apt to cause some temporary damage to turf.

**DIRECTIONS FOR USE: APPLY 20 LBS. PER 2,500 SQ. FT.**

1. Do not apply to a new lawn. Wait until lawn is 3 months old.

58

Lilly's SUPER RICH FEED AND WEED  
Back panel continued ...

12/20/72

39

2. Do not mow lawn for two days prior to application so as to have sufficient weed foliage to receive weed killers.
3. Apply to moist soil and foliage using an accurate spreader and correct dial setting from chart below.
4. Do not water lawn for 24 hours after application but resume watering (if needed) after 24 hours.
5. Make second application after one month if control of hard-to-kill weeds is not complete.

**SPREADER SETTINGS, CALIBRATED TO APPLY 20 LBS. PER 2,500 SQ. FT.**

APPLICATION RATE FOR LILLY'S FEED & WEED IS ALWAYS 20 LBS. PER 2,500 SQ. FT. AS INDICATED BY THESE SPREADER SETTINGS	LILLY SPREADER Model 1652 (White)	LAWN BEAUTY Models 18 & 24	SCOTTS Models 35 & 75
	6½	6	6½

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen N (Minimum) .....12%	Iron as Fe (Minimum) ..... 2%
6% (Minimum) Water Soluble.	(Derived from Sulfasol and Hynite
6% (Minimum) Water Insoluble.	Leather Meal)
(Derived from Urea-formaldehyde,	Sulphur as S (Minimum) ..... 2%
Hynite Leather Meal,	(Derived from Ammonium Sulphate
Ammonium Sulphate and Urea)	and Sulfasol)
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	Magnesium as Mg (Minimum) ..... 1%
(Minimum) ..... 2%	(Derived from Dolomite)
(Derived from Superphosphate)	Zinc as Zn (Minimum) .....0.05%
Soluble Potash (K <sub>2</sub> O) (Minimum) ..... 3%	(Derived from Sulfasol)
(Derived from Muriate of Potash)	Manganese as Mn (Minimum) .....0.05%
Calcium as Ca (Minimum) ..... 2%	(Derived from Sulfasol)
(Derived from Dolomite and Super-	
phosphate)	

**NOTICE**

The Chas. H. Lilly Co. makes no warranty, express or implied, concerning the use of the material, except that it conforms to the chemical description on the label. Neither The Chas. H. Lilly Co. nor the seller shall be held responsible in any way for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

**WARNING**

Induce vomiting if swallowed. Wash hands after use. Keep out of reach of children. In case of contact flush with plenty of water. For eyes get medical attention. Avoid contact with skin, eyes and clothing. Avoid breathing dust. 2,4-D, MCPP and BANVEL-D are injurious to most plants other than grasses so this material should be applied carefully on a day with low wind velocity to avoid drift onto valuable plants. Do not use on St. Augustine grass, Dichondra, carpetgrass or clover lawns. Can cause possible injury to bent grass. Do not apply within the drip line of desirable trees and shrubs. Do not store this material near other insecticides, fungicides, seeds or fertilizers. Rinse spreader with clear water after use. Do not reuse empty container. Destroy it by burying with waste. Dispose of wastes by burying in non-crop lands away from water supplies.

**NOTICE**

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

Manufactured by THE CHAS. H. LILLY CO. Portland, Oregon 97214

59

Copy of label on Printed bag ...  
Front ...

12/20/72



# SUPER RICH

# FEED & WEED

## 12-2-3

ACCEPTED

*Comments in letter*

JAN 19 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 2327-10. SUBJECT  
TO ATTACHED COMMENTS.

*changed to: 802-505*

### HERBICIDE GUARANTEE

#### ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid	.....0.25%
2-(2-methyl-4-chlorophenoxy) propionic acid	.....0.15%
Dimethylamine salt of Dicamba (2-methoxy-3,6-dichlorobenzole acid)	..0.09%

INERT INGREDIENTS .....99.51%

Contains TRIMECTM U.S. Patent No. 3,284,188 T.M.P.B.I. K.C.K.  
R DANVEL-D is the registered trademark of the Velsicol Chemical Co.

**WARNING: KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL CAUTION**

E.P.A. Reg. No. 2327-10-AB-802

## NET WEIGHT 20 LBS.

Back Panel of label on Printed Bag ...

Lilly's Feed & Weed is a real work saver. Through its XF (Extended Feeding) ingredients, it feeds the lawn for weeks, thus reducing the frequency of fertilizing and mowing. TRIMEC, Lilly's effective weed killer, saves the hard work of digging and pulling weeds. Lilly's exclusive PENATURF relieves the surface tension of compact or dense soils to insure that water and food penetrates deep down to the feeding grass roots. This means less watering.

**SUPER RICH FEED & WEED** effectively controls a broad range of lawn weeds, including dandelion, plantain, chickweed, (common & mouse ear), knotweed, clover, Japanese clover, henbit, buttercup, spurge, puncture vine, lawn daisy and other similar weeds.

Apply LILLY'S FEED & WEED any time weeds are growing actively—spring, summer or fall. Applications to moist foliage and soil when temperature is between 50° and 80° F. are the most effective. Applications during hot weather (over 80° F.) or on dry soil are apt to cause some temporary damage to turf.

**DIRECTIONS FOR USE: APPLY 20 LBS. PER 2,500 SQ. FT.**

1. Do not apply to a new lawn. Wait until lawn is 3 months old.

Lilly's SUPER RICH FEED AND WEED  
Back panel continued ...

12/20/72

2. Do not mow lawn for two days prior to application so as to have sufficient weed foliage to receive weed killers.
3. Apply to moist soil and foliage using an accurate spreader and correct dial setting from chart below.
4. Do not water lawn for 24 hours after application but resume watering (if needed) after 24 hours.
5. Make second application after one month if control of hard-to-kill weeds is not complete.

**SPREADER SETTINGS. CALIBRATED TO APPLY 20 LBS. PER 2,500 SQ. FT.**

APPLICATION RATE FOR LILLY'S FEED & WEED IS ALWAYS 20 LBS. PER 2,500 SQ. FT. AS INDICATED BY THESE SPREADER SETTINGS	LILLY SPREADER Model 1662 (White)	LAWN BEAUTY Models 18 & 24	SCOTTS Models 35 & 75
	<b>6 1/2</b>	<b>6</b>	<b>6 1/2</b>

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen N (Minimum) .....12%	Iron as Fe (Minimum) ..... 2%
6% (Minimum) Water Soluble.	(Derived from Sulfasol and Hynite Leather Meal)
6% (Minimum) Water Insoluble.	Sulphur as S (Minimum) ..... 2%
(Derived from Urea-formaldehyde, Hynite Leather Meal, Ammonium Sulphate and Urea)	(Derived from Ammonium Sulphate and Sulfasol)
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) (Minimum) ..... 2%	Magnesium as Mg (Minimum) ..... 1%
(Derived from Superphosphate)	(Derived from Dolomite)
Soluble Potash (K <sub>2</sub> O) (Minimum) ..... 3%	Zinc as Zn (Minimum) .....0.05%
(Derived from Muriate of Potash)	(Derived from Sulfasol)
Calcium as Ca (Minimum) ..... 2%	Manganese as Mn (Minimum) .....0.05%
(Derived from Dolomite and Superphosphate)	(Derived from Sulfasol)

**NOTICE**

The Chas. H. Lilly Co. makes no warranty, express or implied, concerning the use of the material, except that it conforms to the chemical description on the label. Neither The Chas. H. Lilly Co. nor the seller shall be held responsible in any way for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

**WARNING**

Induce vomiting if swallowed. Wash hands after use. Keep out of reach of children. In case of contact flush with plenty of water. For eyes get medical attention. Avoid contact with skin, eyes and clothing. Avoid breathing dust. 2,4-D, MCPP and BANVEL-D are injurious to most plants other than grasses so this material should be applied carefully on a day with low wind velocity to avoid drift onto valuable plants. Do not use on St. Augustine grass, Dichondra, carpetgrass or clover lawns. Can cause possible injury to bent grass. Do not apply within the drip line of desirable trees and shrubs. Do not store this material near other insecticides, fungicides, seeds or fertilizers. Rinse spreader with clear water after use. Do not reuse empty container. Destroy it by burying with waste. Dispose of wastes by burying in non-crop lands away from water supplies.

**NOTICE**

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

Manufactured by **THE CHAS. H. LILLY CO.** Portland, Oregon 97214