JUL \$ 1 1992

Mr. Molly Klettke The Chas. H. Lilly Co. 7737 NE Killingsworth Portland, OR 97318

19 -

Dear Ms. Klettke:

Subject: Revised Labeling Lilly/Miller Go-West Meal EPA Reg. No. 802-507 Your Application Dated June 29, 1992

The labeling submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of the label is enclosed for your records.

Please note that you owe us a response to our letter of June 9, 1992 regarding the active ingredient sodium fluosilicate. A copy of our letter is enclosed.

Sincerely yours,

DHE

Dennis H. Edwards, Jr. Product Manager 19 Insecticide-Rodenticide Branch Registration Division (H-7505C)

74L (4 - 203 - Ref. H

ter sate habiti de contra de la sec

PJK 7

Enclosure

Column: Make and grown hashpround, Milti-ealer Libertrution, also New type. Type sizes: As indicated, all other 6 pt. minimum.

Front panel:

LILLY'S GO WEST MEAL

NEW FORMULA

ACTIVE INGREDIENTS: Sodium Fluosilicate . . . 5.00% Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7tetraoxycyclo-octane) . . . 3.00% INERT INGREDIENTS . . . 92.00% Fluorine Equivalent . . . 3.03%

KILLS SLUGS & SNAILS, Strawberry Root Weevils, Cutworms, Earwigs, Sowbugs

Keep out of reach of children (12 pt. type) AUTION (18 pt. type) This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of trated area. Read and follow all directions. See left side panel for additional precautionary statements.

NET WEIGHT 3 1/2 LBS.

Side and back panels:

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. Avoid contamination of feed and foodstuffs.

IMPORTANT: Keep away from children, dogs and pets. Bait may be ttractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

## STATEMENT OF PRACTICAL TREATMENT

If Inhaled: Remove to fresh air. If affected person is not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention. If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists. If On Skin: Wash with plenty of scap and water.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, swrshes, and potholes). Do not contaminate water when disposing of equipment washwaters. This pesticide may be fatal to dogs or other pets it.

ACCEPTED JUL 31 1992 Vader the Poderel Insected. Parciside, and Roberticitie A ind, for the pest \$202-507 VA Nog. No.

seen. Keep pets out of treated area.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

384

:...

KILLS all these garden pests

Sowbugs Sowbugs Strawberry Root Weevils: Larvae, Adult (illustration) Cutworms Earwigs Snails

READ THE ENTIRE LABEL BEFORE USE, NOTING ALL PRECAUTIONS. Do not use this product on edible feed or food crops.

GO-WEST MEAL is a multi-purpose product designed not only for control of slugs and snails but also strawberry root weevils (larvae and adult), cutworms, earwigs and sowbugs. Since the ingredients are an integral part of the bait and not merely mixed together, GO-WEST MEAL will remain infective longer and will resist washing away by rain and watering.

TO CONTROL LISTED PESTS AROUND ORNAMENTALS, ON LAWNS AND GROUND COVERS

SLUGS AND SNAILS: Since these pests feed on your plants at night and are not generally seen during the day, you may not realize you have an infestation unless you are able to recognize slug damage.

For best results begin applications as early as February and continue baiting into the fall. Apply by broadcast method using 1 lb. GO-WEST MEAL to 250 sq. ft. of area. Scatter bait lightly on ground so that it is barely visible. Do not place in piles.

Re-apply at two-week intervals as long as slugs are present. The aroma of the metaldehyde will attract slugs from a surprising distance. Moist soil increases effectiveness. For effective control-of slug eggs, use filly/Miller Go-West Slug Killer.

EARWIGS - CUTWORMS - STRAWETRRY ROOT WEEVILS - SOWBUGS:

Control these damaging pests by broadcasting GO-WEST MEAL on the ground where insects are seen or damage to plants is noted. Begin baiting for these insects in April.

STORAGE AND DISPOSAL Store this product in its original container and keep in a lbcked . storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash. Filled by weight not by volume. Product may settle after. backaging.

expressed or implied, concerning the effects of this product when not used in accordance with directions. EPA Est. 802-WA-1 EPA Reg. No. 802-507 1505 •• \*\*\*\*\* ••• •• :•..•