

NET WT.
1 1/4 lb.

\$1.98

CAUTION

Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Wash after each use.


Avoid spraying susceptible plants (flowers, shrubs, etc.) Bent grass, Carpetgrass, Dichondra and St. Augustine grass may be injured if contacted. Do not apply within drip line of desirable plants. Do not store near seeds, fertilizers, insecticides or fungicides.

Avoid accidental drift or application to nearby garden plants, flowers, or shrubs. Do not contaminate water used for irrigational or domestic purposes.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-1H

THE CHAS. H. LILLY CO.
PORTLAND, OREGON



Lawn & Turf Weed Bomb

Contains
TRIMEC
Herbicide

Keep from Freezing
Pressurized Spray Ready-to-Use

**SPECIALLY FORMULATED
FOR "HARD-TO-KILL"
LAWN WEEDS**

Contains amines of 2,4-D,
Mecoprop and Dicamba

DIRECTIONS FOR USE

Use to "spot treat" weeds in the lawn. For best results, apply when weeds are actively growing. Do not water within 24 hours after spraying.


Kills...

- | | |
|------------------|------------------|
| Dandelions | Canada Thistle |
| Plantains | Filaree |
| Chickweed | Henbit |
| Clover | Lambsquarters |
| Black Medic | Mustard |
| (Yellow or | Pennywort |
| Japanese Clover) | Pigweed |
| English Daisy | Sheep Sorrel |
| Curly Dock | Yarrow |
| Heal-All | Buttercup |
| Ground Ivy | Oxalis |
| Knotweed | Shepherd's Purse |

Shake container well before using. Miller's Lawn and Turf Weed Bomb is designed to permit easy treatment of lawn weeds without bending or stooping. Invert can, aim at weed and press valve. Only a short burst is needed. Over-spraying only wastes the spray.

Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 3-4 weeks if necessary.

For lawns with many weeds, requiring an overall treatment, spray with Miller's Lawn & Turf Weed Killer.



CAUTION Keep out of the reach of children
See left side panel for additional cautions.
E.P.A. Reg. No. 802-513-AA

NET WT.
1 1/4 lb.

5/98

CAUTION


Avoid contact with eyes. Wash thoroughly if contact occurs. Avoid contact with clothing. In case of contact flush with plenty of water. If eyes get medical attention. Wash with water.

Avoid spraying susceptible plants. Flowers, shrubs, etc. Best used on Carpetgrass, Dichardra and St. Augustine grass. May be used if contacted. Do not apply within 100 ft. line of desirable plants. Do not use near seeds, fertilizer, herbicides or fungicides.

Avoid accidents. Drift or application to desirable plants, flowers, shrubs. Do not contaminate water supply. Do not get material on clothing.

WARNING: Contents under pressure. Do not puncture. Do not use near fire or heat. Do not use in excessive temperatures. For use only in the United States and Canada. Do not use in other countries.

© 1988 F. S. I. Garden Corp.
All rights reserved.
Printed in the U.S.A.



Lawn & Turf Weed Bomb



Active Ingredients:

- *Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 0.130%
- **Dimethylamine salt of 2,4-dichlorophenoxyacetic acid . 0.267%
- ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) 0.029%

Inert ingredients 99.554%

*Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid 0.100%

**Equivalent to 2,4-dichlorophenoxyacetic acid 0.200%

***Equivalent to 3,6-dichloro-o-anisic acid 0.024%

*TRIMEC® is a Reg. T.M. of F.S.I.-Garden Corp. U.S. Patent No. 3,284,166

DIRECTIONS FOR USE

Use on lawn and turf areas. For best results, apply when weeds are actively growing. Do not water within 24 hours of treatment.

Kills...

- | | |
|-----------------|-----------------|
| Crabgrass | Spina-Top® |
| Field Bindweed | St. John's Wort |
| Flower Spurge | St. Louis Lily |
| Goosegrass | Summer Crocus |
| Groundsweeper | Summer Squash |
| Jack-in-the-box | Summer Squash |
| Yellow Pine | Summer Squash |
| Yellow Top | Summer Squash |
| English Daisy | Summer Squash |
| Blue Dandelion | Summer Squash |
| Red Top | Summer Squash |
| Red Top | Summer Squash |
| Red Top | Summer Squash |

Use on lawn and turf areas. For best results, apply when weeds are actively growing. Do not water within 24 hours of treatment.

Use on lawn and turf areas. For best results, apply when weeds are actively growing. Do not water within 24 hours of treatment.

Use on lawn and turf areas. For best results, apply when weeds are actively growing. Do not water within 24 hours of treatment.

CAUTION Keep out of the reach of children. Do not use on lawns or other areas where children play.