

1/15/62 516

1/15/62

DIRECTIONS FOR USE

Use 2 tablespoons Lily Miller Vegetable and Fruit Spray per gallon of water. Apply spray when insects first appear and repeat as necessary to maintain effective control. Coverage of both surfaces of leaves is essential for good control. Observe use limitations and warnings. Observe days interval between last spraying and harvest as indicated by number shown in parenthesis following each crop.

VEGETABLES

Apply 1 gallon mixed spray to 275 sq. ft. of garden area.

BEANS, SUCCULENT (Green Beans) (3) — To control Black Bean Aphid, Mexican Bean Beetle. (Do not apply to Lima Beans. Do not apply this product more than 3 times per season. Do not feed treated forage or freshings to livestock. Do not graze livestock in treated gardens or fields.)

BROCCOLI (7), BRUSSELS SPROUTS (14), CABBAGE (7), CAULIFLOWER (14). To control Cabbage Aphid, Cabbage Looper, Imported Cabbageworm, Diamond Back Moth Larvae, Flea Beetle, Marquon Bug.

CUCUMBERS (0), MELONS (0), SQUASH Summer and Winter (0) — To control Aphids, Cucumber Beetle, Squash Bug, Striped Flea Beetle.

PEPPERS (4) — To control Green Peach Aphids, Flea Beetle, Pepper Maggot, Hornworms.

POTATOES (0) — To control Green Peach Aphid, Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper, Whiteflies, Armyworm.

LETTUCE, HEAD (14) — To control Green Peach Aphid, Cabbage Looper, Diamond Back Moth Larvae, Imported Cabbageworm. (Do not apply more than 3 times after thinning. Remove outside leaves at harvest. Do not feed crop waste to livestock.)

TOMATOES (1) — To control Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Whiteflies, Tomato Hornworm, Tomato Russet Mite, Yellow-striped Armyworm.

FRUIT and NUTS

APPLES (30) — To control Apple Aphid, Romy Apple Aphid, Apple Rust Mite. (Apply up to 1 gallon mixed spray per tree. Do not apply more than 3 times during fruiting period. Do not feed pomace from treated apples to livestock.)

APRICOTS (30), PEACHES (30), NECTARINES (30) — To control Caterfacing Insects, Green Peach Aphid, Peach Twig Borer, Peach Silver Mite. (Apply 1 to 2 gallons mixed spray per tree. Do not apply more than twice during fruiting season.)

PEACHES (21), PRUNES (7) in OREGON — Western Peach Tree Borer. (Root Borer). (Makes first application during period of July 5 to 15th and a second one in three weeks. Spray trunk and lower 4' of limbs. Let spray puddle around base. Avoid spraying fruit. Do not make more than 2 applications when fruit is present.)

PEARS (7) — To control Pear Leaf Blister Mite, Pear Rust Mite. (Apply up to 1 gallon of mixed spray per tree. For Pear Leaf Blister Mite, make one application during post-harvest or dormant season. Do not make more than two applications during fruiting season.)

Continued on right panel.



Vegetable & Fruit Spray



Containing Thiodan* For Control of Certain Insects on Vegetables, Fruits and Ornamentals

WARNING Keep out of reach of children. See right side panel for additional warnings.

Continued from left panel.

PEARS (21) — To control Pear Psylla. (Use up to 1 1/2 gallons mixed spray per tree. Begin applications when nymphs are small or when adults are first noticed. Repeat sprays as necessary, but do not exceed 5 applications during fruiting season.)

STRAWBERRIES (4) — To control Strawberry Aphids, Cyclamen Mites, Meadow Spittlebug. (Apply 1 gallon mixed spray to 275 sq. ft. of garden area. Do not reapply within 15 days or more than twice within a 35-day period when fruit is present.)

WALNUTS — To control Walnut Aphid. (Apply 1-1 1/2 gallons mixed spray per tree. Do not apply after husk split. Do not graze livestock in treated area.)

ORNAMENTALS, Out-of-doors — To control Aphids, Cyclamen Mites, Whiteflies, Rose Chafers. Use 2 tablespoons per gallon of water. Do not use on Birch Trees. On Chrysanthemums, best results will be obtained if applied before plants flower. Do not apply to "Bonnie DeLuxe", "Fred Shoemith" and "White Knight" chrysanthemums as injury may result. NOTE: Birds feeding on treated areas may be harmed.

WARNING

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing.

Note to Physicians: Thiodan is a central nervous system stimulant. Treat symptomatically. There is no specific antidote.

During commercial or prolonged exposure in spray mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Bureau of Mines for Thiodan protection.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not allow spray to drift onto neighboring gardens or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply when runoff is likely to occur. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Destroy empty container. Do not re-use for any purpose. Do not use or store near heat or open flame.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

LC-2N

EPA Reg No. 802-516

EPA Est. 802-OR-1

BEST AVAILABLE COPY

ACCEPTED
5 1979
EPA Reg No. 802-516