

SPECIFIC DIRECTIONS

TANSY RAGWORT CONTROL (in Pastures, along Fence Rows, Drainage Ditchbanks, Roadsides, Rangelands)

1. Use 3 qts. in 50 gallons of water per acre. On small areas, use 4 tablespoons in 1 gallon of water.
2. Apply in the spring before any flowers appear (during rosette stage of growth). The earlier the application in relation to plant growth, the better the kill. Make a thorough spray application to wet all the foliage thoroughly.
3. Respraying for more than one year is necessary to control late germinating seeds. May be applied in early fall after beginning of regrowth (crowns) or newly germinated seeds.
4. Do not graze dairy animals on treated areas within 7 days after application.

PASTURE WEED CONTROL . . . Dandelion, Plantain, Bur Clover

1. Use 1½ to 2 pints per acre in sufficient water to give uniform coverage.
2. **CAUTION** — Do not apply on Creeping Grasses, such as Bent, except for spot spraying. Do not apply on newly seeded pastures until grass is well established. Most legumes are usually damaged or killed. Do not apply to Dichondra or St. Augustine grasses.
3. Do not graze dairy animals on treated areas within 7 days after application.

WOODY PLANT CONTROL (Rangeland, pastures, along fence rows, drainage ditchbanks, roadsides, under power lines, and other similar waste areas)

1. Use 3 qts. per 100 gal. of water for such woody plants as Willow, Elderberry, Sumac, Wild Cherry, Wild Grape.
2. Apply as a drenching spray for brush up to 8 feet tall, after foliage is well developed.
3. Make thorough coverage of all parts of the plant, including foliage, shoot stems, and bark.
4. Do not graze dairy animals on treated areas within 7 days after application.



LV 2,4-D ESTER FOUR BP

(4 lbs. 2,4-D acid equivalent per gallon)

For the control of many Broadleaved Weeds and certain Woody Plants

Dandelion	Peppergrass	Tarweed	Wild Morning
Docks	Pigweed	Thistles	Glory,
Fanweed	Plantains	Vetches	and
Lamb's Quarters	Ragweed	White Top	Many other
Mallow	Shepherd's Purse	Wild Carrot	broadleaved
Mustards	St. Johnswort	Wild Radish	weeds

Also for . . . Alder, Elderberry, Hazel, Coastal Sage, Sand Sagebrush, Sumac, Willow, Buckbrush

A LOW VOLATILE 2,4-D HERBICIDE

ACTIVE INGREDIENT:	
2,4-Dichlorophenoxyacetic Acid, Butoxypropyl Esters	72.52%
INERT INGREDIENTS:	27.48%
Total	100.0%
(2,4-D Acid Equivalent 44.3%)	

E.P.A. Reg. No. 802-517-AA

WEED KILLER

CAUTION — Keep out of reach of children

See right side panel for additional cautions

NET CONTENTS: _____ GALLON(S)

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

Miller's

CAUTION

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.

Harmful if swallowed

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water, flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children and domestic animals.

Do not use around the home or recreation area.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, can berries, ornamentals or other desirable plants which are sensitive to 2,4-D and do not permit spray mist to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Application by airplane, ground equipment, or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4-D susceptible plants.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

Ester formulations may cause more crop injury than amine formulations. Do not use on crops other than those listed on this label.

Be sure that the use of this product conforms to all applicable local regulations.

At higher temperatures vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.