

LV 2,4,5-T ESTER FOUR BP

GENERAL DIRECTIONS

Add half the required amount of water or oil to the spray tank, then add the specified quantity of this product with agitation, then add balance of the water or oil with continued agitation.

NOTE...If this product is to be used in straight oil mixtures, do not let water get into the finished mixture. The product forms an emulsion in water - not a solution. The water emulsion tends to separate on standing and for this reason continued agitation is necessary to prevent separation and to maintain a uniform spray mixture.

SPECIFIC DIRECTIONS

BRUSH FOLIAGE SPRAY

- 1. For Fence Rows, and Right-of Ways, use 3 to 4 qts. per 100 gals.of water.
- 2. For **Pastures**, apply 3 qts. per acre when grass is well established. Do not apply to newly seeded areas. For **Rangelands**, apply 4 qts. per acre in the spring when brush is in heavy foliage stage (40 to 90 days after leaves unfold). Use in sufficient water (100 gals or more by ground equipment) to give complete coverage of all brush.
- 3. On woody plants, ranging from 5 to 6 feet tall, treatments should be made when plants are actively growing and the leaves are fully developed. Apply with a power or hand sprayer with nozzle adjusted to give coarse spray
- 4. Coverage should be complete. All parts of the plants, including foliage, shoot stems and bark should be wet with the spray.
- 5. If new growth appears, repeat application.

STUMP AND BRUSH STUB SPRAY

- 1. Use 3 to 4 gts. in 25 gals of fuel oil or kerosene. Mix thoroughly.
- 2. Cut growth over 5 to 6 feet tall near the ground, then spray freshly cut stumps. Be sure to cover thoroughly bark and cut surfaces.
- 3. If sprouts 3 or 4 feet tall are present from previously cut stump, spray thoroughly using 3 qts. per 100 gallons of water.

SPECIFIC ADVICE

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

Consult State Agricultural Experiment Station or State Extension Service Weed Specialists for specific advice as climate and local conditions may cause suggested uses to vary.

NOTICE: Follow directions carefully. Fining and method of application, weather and insprace directors in the weather as a specifically recommended, and other influencing factors in the use of this product are beyond the instrument of the sector. Buyer assume a risks of use storage or tandling of this material of the storage of tandling of the storage of tandling of the storage of the storage of tandling of the storage of the storage of tandling of the storage of the storage of tandling of tandling of the storage of tandling of tandlin

LV

For the control of Herbaceous and Woody Plants growing in Pastures, Range Land, Fence Rows, **Manual Science** and Right-of-Ways.

.Ash

C

Horse Nettle



LV 2,4,5 -T ESTER FOUR BP

Brambles

Chokecherry

Hawthorn

Maple

Oak

Poison Oak

Poison Ivy

Salmonberry

Wild Blackberry

Wild Cherry

Wild Rose, and

Certain Species

of Ribes

9nsist on ...



LV 2,4,5-T E FOUR I

(4 lbs. 2,4,5-T acid equivalent

For the control of Herbaceous and Wood Pastures, Rangeland, Fence Rows

A LOW VOLATILE 2,4,5-T

ACTIVE INGREDIENT *2,4.5 Trich prophenoxyacety, and Butoxypropy, Esters INERT INGREDIENTS

• • Fall Later 1 : 44 7 - 24 F 7 A

EPA Reg. No. 802-523-A

WEED KILLEF

CAUTION --- Keep out of reach See right side panel for additional caution

NET CONTENTS:

1 . .

THE CHAS. H. LILLY C PORTLAND, OREGON EPA Est 802 OR 1

Miller's

LV 2,4,5-T **ESTER** FOUR BP

For the control of Herbaceous and Woody Plants growing in Pastures, Range Land, Fence Rows, and Right-of-Ways.

Ash

Brambles Chokecherry Hawthorn Horse Nettle Maple Oak **Poison Oak** Poison Ivy Salmonberry Wild Blackberry Wild Cherry Wild Rose, and Certain Species of Ribes



LV 2,4,5-T ESTER FOUR BP

(4 lbs. 2,4,5-T acid equivalent per gallon)

For the control of Herbaceous and Woody Plants growing in Pastures, Rangeland, Ferice Rows and Right-of-Ways

A LOW VOLATILE 2,4,5-T HERBICIDE

ACTIVE INGREDIENT *2,4,5-Trichlorophenoxyacetic acid, Butoxypropyl Esters INERT INGREDIENTS

*Equivalent to 44.7% 2,4,5 T. Acid.

E.P.A. Reg. No. 802-523-AA



CAUTION — Keep out of reach of children See right side panel for additional cautions

NET CONTENTS:

THE CHAS. H. LILLY CO PORTLAND, OREGON EPA Est 802 OR-1

the or

get ion. ued bray

100 not the

bld). give

ade pply ray. 100t

pray

cut

tion. hter.

Vecd sted

cals not mes all MP 1 1





68.6% 31.4%

GALLON(S)



CAUTION

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.

Harmful if swallowed.

This product may cause skin irritation Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water, flush eyes with plenty of water for at least 15 minutes and get medical attention Keep out of reach of children and domestic animals.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, cane berries, ornamentals or other desirable plants which are sensitive to 2,4,5-T and do not permit spray mist to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Clovers may be injured with this material Do not use on creeping grasses such as Bent Grass and St. Augustine Grass.

Application by airplane, ground equipment, or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4,5-T susceptible plants.

Do not contaminate irrigation ditches or water used for irrigation or domestic purposes. Excessive amounts of 2,4,5-T in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not burn.

Do not use around the home or recreation areas.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.