

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the products registered under EPA Reg. No. 802-525

Avoid excessive wetting of blossoms. A light application is all that is necessary. Apply just a short burst of spray. Avoid soaking the flowers, leaves or stems as this may cause imperfectly formed fruit. Seedless tomatoes are produced by spraying first blossoms as soon as they open before pollination occurs, otherwise fruits will have seed.

CAUTION

Do not spray near open flame or near sparks. Do not smoke while spraying. Avoid exposure of container to high temperatures (120°F). Store this container away from stoves, radiators, other heat sources or direct sunlight.

**WARNING
CONTENTS UNDER PRESSURE**

Never set container on stoves, radiators, nor place where temperature will exceed 120° F and may cause it to burst. Do not puncture or throw container into fire.

ACTIVE INGREDIENT:
* p-chlorophenoxyacetic acid 0.0050%
INERT INGREDIENTS: 99.9950%
*Equivalent to 50 ppm

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



MP-1K

Manufactured for
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 10900 CA 1

**NET WT.
5 1/2 OZ.**



**Super
Tomato
set**



CAUTION Keep out of reach of children.
See left side panel for additional statements.

During the early part of the season, cloudy days, cool nights and short periods of sunlight can cause poor pollination of the first tomato flowers, resulting in fewer fruit being set, or fruit not set early enough to ripen properly.

Miller's Super Tomato Set aids the early tomato flowers to set fruit even under adverse weather conditions.

DIRECTIONS FOR USE

1. Hold sprayer at least 10 to 15 inches from blossoms.
2. Earliest tomatoes are secured by making a spot application to the first cluster when most of the flowers in the cluster are open before pollination.
3. For **general application**, direct spray nozzle at blossom cluster and apply a short burst of spray, just enough to moisten blossoms. Avoid wetting the foliage as much as possible. Release spray valve when moving from one blossom cluster to the next so as not to wet foliage in between.
4. Repeat spray at 7 to 10 day intervals to set blossoms opening later but do not make more than 5 applications during the season and do not apply more than once to individual flower clusters. Do not apply within 15 days of harvest.

USE PRECAUTIONS

Do not spray the small flower buds that are just developing. This may cause them to drop before setting fruit. Do not hold can too close to blossoms or spray them too long as some temporary damage to foliage or buds may occur. This can be avoided by holding spray 10 to 15 inches from blossoms, giving them a short burst of spray and timing spray application as above.