

802530

ACCEPTED
FEB 22 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 802530

LILLY/MILLER
CHLORBAN
INSECT SPRAY
Contains DURSBAN* Insecticide

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-diethyl 0-(3, 5, 6-trichloro-2-pyridyl)
phosphorodithioate) 6.79%

INERT INGREDIENTS. 93.21%


*Dursban® Reg. TM of the Dow
Chemical Co. U.S. Pat. No. 3,244,586

IDEAL ANT CONTROL

Also controls Chinch Bugs, Earwigs,
Fleas, Cutworms, Sod Webworm
(Lawn Moth Larvae) in lawns
Aphids, Mites, Spittlebugs,
Whiteflies on Ornamentals

Keep out of reach of children
WARNING
See left side panel for Statement of Practical
Treatment and additional Precautionary Statements

NET CONTENTS _____ (1 Pint or 1 Quart)



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through skin. May be injurious to eyes and skin. Do not take internally. Avoid breathing spray mist or vapors. Wash thoroughly after handling or after contact with skin. Do not smoke while using. Keep children and pets off treated areas until spray has dried. Handle concentrate in a ventilated area. Do not spray vegetable gardens or other plants used for food purposes, except as specifically directed.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes, flush with plenty of water and get medical attention. If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARD
DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USING

LAWNS AND TURF To control Billbugs, Chinch Bugs, Sod Webworm (Lawn Moth Larvae), Ants, Brown Dog Ticks, Fleas, Clover Mites, Armyworms, Crickets, Cutworms, Earwigs and Grasshoppers infesting home lawns and ornamental or recreational turf grass areas, mix Lilly/Miller Lawn Insect Spray with water according to the directions given in the following table and apply as a coarse droplet spray using a low pressure tank sprayer, or a hose attachment sprayer (such as Lilly/Miller Big 15). For best results the lawn should be moist at the time of application and care should be taken to obtain complete and uniform coverage of infested areas. Spray when pests first appear and re-treat when needed,

DOSAGE AND DILUTIONS FOR USE ON LAWNS AND TURF

Area to be treated	Amount of Lilly/Miller Chlorban Insect Spray	Dilute with water to make
100 sq. ft.	3½ teaspoons	3 gallons
500 sq. ft.	6 tablespoons	15 gallons
1000 sq. ft.	12 tablespoons (6 fl. ounces)	30 gallons
1500 sq. ft.	18 tablespoons (9 fl. ounces)	45 gallons

FOR CONTROL OF EUROPEAN CRANE FLY LARVAE (LEATHER JACKETS)

Area to be treated	Amount of Lilly/Miller Chlorban Insect Spray	Dilute with water to make
500 sq. ft.	9 tablespoons (4½ fl. ounces)	15 gallons
1000 sq. ft.	18 tablespoons (9 fl. ounces)	30 gallons
1500 sq. ft.	27 tablespoons (13½ fl. ounces)	45 gallons

FOR CONTROL OF BILLBUGS

Area to be treated	Amount of Lilly/Miller Chlorban Insect Spray	Dilute with water to make
500 sq. ft.	¾ - 1½ cups	(6 - 12 fl. oz.) 15 gallons
1000 sq. ft.	1½ - 3 cups	(12 - 24 fl. oz.) 30 gallons
1500 sq. ft.	2¼ - 4½ cups	(18 - 36 fl. oz.) 45 gallons

NOTE: To help prevent infestation of buildings by Ants, Crickets, Earwigs and Sowbugs (pillbugs), treat a band of soil 6 to 10 feet around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

ORNAMENTAL PLANTS On flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with Aphids, Bagworms, Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Fall Webworm, Spittlebugs, Thrips (exposed) and Whiteflies, use Lilly/Miller Lawn Insect Spray according to the following table and apply with a suitable pressure sprayer or a hose attachment sprayer (such as Lilly/Miller Six-Shooter). Make certain of complete, uniform coverage of infested plants. Avoid over-spraying to the point of excessive run-off. Treat when pests appear and repeat at 7 to 10 day intervals if needed.

IMPORTANT: Do not use on Azaleas, Camellias, Poinsettias, Roses, or variegated Ivy because of possible injury to these plants.

DOSAGE TABLE FOR ORNAMENTAL PLANTS

Pest Controlled	Amount of Lilly/Miller Chloroban Insect Spray in water to make	
	1 gallon	3 gallons
Aphids Mites	4 teaspoons	4 tablespoons
Bagworms Spittlebugs		
Fall Webworm Whiteflies		
Grasshoppers		
Cutworms Leafhoppers	8 teaspoons	½ cup (8 tablespoons)
Mealybugs Thrips (exposed)		

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental insects will be killed for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.

NOTE: To help prevent infestation of buildings by Ants, Crickets, Earwigs and Sowbugs (pillbugs), treat a band of soil 6 to 10 feet around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.


ORNAMENTAL PLANTS On flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with Aphids, Bagworms, Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Fall Webworm, Spittlebugs, Thrips (exposed) and Whiteflies, use Lilly/Miller Lawn Insect Spray according to the following table and apply with a suitable pressure sprayer or a hose attachment sprayer (such as Lilly/Miller Six-Shooter). Make certain of complete, uniform coverage of infested plants. Avoid over-spraying to the point of excessive run-off. Treat when pests appear and repeat at 7 to 10 day intervals if needed.

IMPORTANT: Do not use on Azaleas, Camellias, Poinsettias, Roses, or variegated Ivy because of possible injury to these plants.

DOSAGE TABLE FOR ORNAMENTAL PLANTS

Pest Controlled		Amount of Lilly/Miller Chlorban Insect Spray in water to make	
		1 gallon	3 gallons
Aphids	Mites		
Bagworms	Spittlebugs	4 teaspoons	4 tablespoons
Fall Webworm	Whiteflies		
Grasshoppers			
Cutworms	Leafhoppers		
Mealybugs	Thrips (exposed)	8 teaspoons	½ cup (8 tablespoons)

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental insects will be killed for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.



FOR CONTROL OF ROOT MAGGOTS ON CERTAIN DIRECT-SEEDED VEGETABLE CROPS

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage:

For direct seeded crops use 2 to 4 tablespoons in a gallon of water and apply with sprinkler can in a band 2 to 4 inches wide over 100 linear feet of row at planting time.

Radishes:

Use 3½ teaspoons in a gallon of water and apply as a soil drench in the seed furrow, with the seed, to 100 linear feet of row at planting time.


Restrictions: Do not make more than one application per planting.

Rutabagas, Turnips:

Use 2 to 4 tablespoons in a gallon of water and apply in a band 2 to 4 inches wide over a 100 linear feet of row at time of planting. With rutabagas it would be realistic to only expect partial protection because of their extended growing season.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Rinse thoroughly before discarding and securely wrap original container in several layers of newspaper and discard in trash.



NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

LC-5Q

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

EPA Reg. No. 802-530

EPA Est. 802-OR-1

