

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.


WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonable fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

**THE ANDERSONS
LAWN FERTILIZER DIVISION
P.O. BOX 119
MAUMEE, OH 43537**



SEP 12 1990

Ms. Molly Klettke
The Chas. H. Lilly Co.
7737 NE Killingsworth
Portland, OR 97218

Dear Ms. Klettke:

Subject: Added use
Lilly/Miller Chlorban Insect Granules
EPA Registration No. 802-532
Amendment application of November 21, 1989

The Amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below:

Delete the claim for control of cockroaches around homes and buildings
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If These conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

275

-2-

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Dennis H. Edwards, Jr.
Product manager (12)
Insecticide-Rodenticide Branch
Registration Division (H-7505C)

Enclosure

LILLY/MILLER
CHLORBAN INSECT GRANULES

ACTIVE INGREDIENT:

Chlorpyrifos (0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate) 0.5%
INERT INGREDIENTS 99.5%

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 12 1990

DURSBAN is a trademark of The Dow Chemical Company

Controls Root Maggots in certain vegetable crops;
Ants, Armyworms, Grasshoppers and many others in lawns.

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

802-532

Keep out of reach of children

CAUTION

See left side panel for Statement of Practical Treatment and additional
Precautionary Statements.

NET WEIGHT 3 LB.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep out of reach of children. May be harmful if swallowed.
Do not take internally. Avoid eye contact. Wash thoroughly after
handling. Wash contaminated clothing before reuse. Keep children and
pets off treated areas until this material is washed into the soil and
treated area is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2
glasses of water and induce vomiting by touching back of throat with
finger. Do not induce vomiting or give anything by mouth to an
unconscious person. If Inhaled: Remove victim to fresh air if symptoms
of cholinesterase inhibition appear and get medical attention
immediately. If On Skin: Flush with plenty of water. If In Eyes: Flush
with plenty of water. Get medical attention. NOTE TO PHYSICIAN:
Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically.
Atropine, by injection only, is antidotal.

ENVIRONMENTAL HAZARDS


This product is toxic to fish, birds and other wildlife. Keep out of
any body of water. Do not contaminate water when disposing of equipment
washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

FOR LAWNS AND OTHER ORNAMENTAL TURF AREAS

To control Ants (including foraging fire ants), Armyworms, Billbugs,
Centipedes, Chiggers, Chinch Bugs, Crickets, Cutworms, Earwigs, Fiery
Skipper Larvae, Flea Beetles, Fleas, Gnats, Grasshoppers, Greenbugs,
June Beetles, Leafhoppers, Lucerne Moth, Millipedes, Mites (Clover,



Formula Grass, Winter Grain), Mosquitoes, Sod Webworms (Lawn Moth Larvae), Sowbugs (Pillbugs), Spittlebugs, Springtails, Ticks, and Tropical Grass Webworms.

Apply CHLORBAN INSECT GRANULES at the rate of 1 lb. per 200 sq. ft. of area.

Treat when insects first appear. Use any hand operated gravity spreader designed for application of granular materials. Immediately after application, sprinkle the turf with 6 gallons of water per 200 sq. ft. so as to wash the granules into the zone of insect infestation.

WHITE GRUB CONTROL (Larvae of European Chafer and Japanese Beetle)
Apply 1 lb. CHLORBAN INSECT GRANULES per 50 - 100 sq. ft. Apply the granules uniformly over the area to be treated, followed immediately by a thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service for proper time to treat in your area.

EUROPEAN CRANE FLY LARVAE CONTROL
Apply 1 lb. CHLORBAN INSECT GRANULES per 140 sq. ft. Use any hand operated gravity or rotary spreader designed for application of granular materials. Immediately after application, sprinkle the turf with 3 gallons of water per 100 sq. ft. For timing of applications you should consult your local Agricultural Extension Service.

HYPERODES (Turfgrass) WEEVIL CONTROL
Apply CHLORBAN INSECT GRANULES at the rate of 1 lb. per 100 sq. ft. of area. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service Specialist for best time to treat in your area.

NUISANCE PESTS AROUND HOMES AND BUILDINGS - OUTDOORS
To help prevent infestation of homes and buildings by Ants (including foraging fire ants), Bees, Boxelder Bugs, Carpenter Ants, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Hornets, Millepedes, Mosquitoes, Scorpions, Silverfish, Sowbugs (Pillbugs), Spiders, Ticks, Wasps and Yellowjackets:

Treat a band of soil 6 to 10 feet around and adjacent to the building where pests are active and may find entrance. Apply CHLORBAN INSECT GRANULES at the rate of 1 lb. per 200 sq. ft. (10' x 20'). Use any hand operated gravity spreader designed for application of granular materials. Mosquitoes coming to rest on treated areas will be killed for varying periods of time. For Scorpions, remove debris such as lumber, firewood and other materials which serve as shelter for these insects.

FOR ROOT MAGGOT CONTROL
IN DIRECT SEEDED CROPS: Apply only at seeding. Do not make more than one application per season.

