

LILLY/MILLER
CASORON GRANULES

ACTIVE INGREDIENT:

Dichlobenil* 2.0%
INERT INGREDIENTS 98.0%

Trademark and research discovery of Duphar B.V. Amsterdam, Holland.

*2,6 Dichlorobenzonitrile


CONTROLS WEEDS & GRASSES
IN EVERGREENS, ROSES, SHRUBS, FRUITS & NUTS

TREATS 270 SQ. FT.

ACCEPTED
MAR 28 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 802-536

Keep out of reach of children
CAUTION
See right side panel for additional cautions.

NET WT. 1 1/4 LBS.



Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not breathe dust. Do not allow contact with eyes or skin. In case of contact, flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Avoid contamination of feed and foodstuff.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Casoron Granules will selectively control annual grassy and certain perennial weeds around established ornamental trees and shrubs, fruit and nut trees, and berries, listed on this label. One application, made as directed, is generally enough to give season-long control.

CASORON GRANULES when applied to the soil as directed will give selective control of annual grassy and broadleaf weeds and certain perennial weeds such as:

Annual Bluegrass	Field Morning Glory	Ragweed
Artemesia	Florida Purslane (Pusley)	Red Deadnettle
Bluegrass	Foxtail	Redroot Pigweed
Bull Thistle	Gisekia	Rosarypea
Camphorweed	Groundsel	Russian Knapweed
Canada Thistle	Henbit	Russian Thistle
Carpetweed	Horsetail	Shepherdspurse
Chickweed	Jerusalem oak Goosefoot	Smartweed
Citron Melon	Knotweed	Spanishneedles
Coffeeweed	Lambsquarters	Spurge
Crabgrass	Milkweed Vine (Morrenia odorata)	Teaweed
Cudweed	Natalgrass	Texas Panicum (Hurrahgrass)
Curly Dock	Peppergrass	Wild Carrot
Dandelion	Pineapple Weed	Wild Mustard
Dogfennel	Plantain	Wild Radish
Eveningprimrose	Purslane	Yellow Woodsorrel
False Dandelion	Quackgrass (from seed)	
Fiddleneck		

BEST AVAILABLE COPY

CASORON GRANULES are to be applied only to established woody plants – those which have been planted for at least four months.

CASORON GRANULES may be used for weed control on the following Ornamental Woody Shrubs:

Amur Corktree	Euonymous	Mock Orange
Arborvitae	Filbert	Mountain Ash
Ash	Flowering Almond	Nandina
Azalea 1	Flowering Quince	Oak
Barberry	Forsythia	Osmanthus (Holly-O)
Beauty Bush	Gardenia	Pachistima
Birch	Goldenraintree	Pecan
Box Eider	Hackberry	Photinia
Boxwood	Heather	Pittosporum
Camellia	Holly 2	Poplar
Caragana	Honeysuckle	Privet
Cleyera Japonica	Ivy (Hedera Helix)	Pyracantha
Cotoneaster	Juniper	Rhododendron
Cottonwood	Kinnikinnick	Rockrose
Crabapple	Laurel	Rose
Deutzia	Leucothoe	Russian Olive
Dogwood	Lilac	Spiraea
Eastern Red Cedar	Linden	Squaw-carpet
Elm	Locust	Weigela
Elwood Cypress	Magnolia	Willow
English Walnut	Maple	Yew (Taxus)

- 1 Kurume, Mollis, Hardy Hybrid types and hardy native species only.
- 2 Except I. Crenata, I. Rotunda, I. Vomitoria.

CASORON GRANULES may be used for weed control on the following Edible Fruits & Nuts:

Almond	Filbert	Nectarine
Apple	Grape	Orange
Blackberry	Grapefruit	Peach
Blueberry	Lemon	Pear
Cherry	Lime	Prune-Plum
English Walnut		Raspberry

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Before Application

1. Remove existing weeds.
2. Thoroughly work and smooth the area to be treated.
3. Carefully measure area to be treated and determine the number of square feet.
4. If area to be treated is large, it should be divided into smaller sections and each treated individually.

Application and Rate

1. Shake Casoron Granules uniformly on the soil at the rates specified.

ANNUAL WEEDS: Apply in early spring before weeds emerge or after removing weeds.

PERENNIAL WEEDS: For best control, apply after a killing frost from late fall to early winter.

2. FRUIT TREES, NUTS, BLUEBERRIES, GRAPES: Apply 2-3 level Tbs. per 10 sq. ft. 20 oz. treats 180-270 sq. ft. Use the higher rate- 3 tablespoons per 10 sq. ft. or 20 oz per 180 sq. ft. - if weed infestation is heavy or if most weeds are perennial. Wait 30 days after making application before harvesting stone fruits or nuts.

BLACKBERRIES, RASPBERRIES- Use 2 level tablespoons per 10 sq.ft. Use only on established plantings. Apply before new shoots emerge in spring.

CITRUS: Use $1\frac{1}{2}$ to 2 tablespoons per 10 sq. ft. area. Apply around trees which have been planted for at least one year. Use higher rate for heavy weed infestation or if most weeds are perennial.

ORNAMENTAL WOODY SHRUBS: Use at rate of $\frac{3}{4}$ ounce (2 level tablespoons) for each 10 sq. ft. to be treated. 20 oz. treats 270 sq. ft.

3. Use the diagram below as a guide for uniform distribution of granules.

DIAGRAM

4. Rates higher than those indicated may cause injury.
5. More accurate and uniform application can be obtained by applying one-half of the required material over the entire treatment area and repeating the application with the remaining material.
6. CASORON GRANULES should not be used on sloping areas, because run-off may result in injury to lawn or other plants in adjacent areas.

After application

1. Unless the soil is wet, sprinkle lightly to moisten the soil $\frac{1}{4}$ to $\frac{1}{2}$ inch deep. Do not flood, avoid puddling and washing.
2. Avoid disturbing soil in the treated area as long as weed control is being maintained.



STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals.

Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

NOTE: Use only on the plants listed. Do not apply to transplant or cutting beds or to newly set transplants, and do not seed or transplant into treated areas for one year after application. Do not use on extremely light sandy soils.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

29802

THE CHAS. H. LILLY CO.
Portland, Oregon 97218

EPA Est. 802-OR-1

EPA Reg. No. 802-536

