

LILLY/MILLER  
Casoron Granules

Controls Weeds & Grasses in Evergreens, Roses, Shrubs, Fruits & Nuts

ACTIVE INGREDIENT:

Dichlobenil: 2,6-Dichlorobenzonitrile . . . . 2.0%

INERT INGREDIENTS . . . . 98.0%

Trademark and research discovery of Duphar, B.V., Amsterdam, Holland.


TREATS 2,133 SQ. FT.

Keep out of reach of children

CAUTION

See back panel for additional cautions

NET WEIGHT 10 LBS.



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Do not allow contact with eyes or skin. In case of contact, flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Avoid contamination of feed and foodstuff.

ENVIRONMENTAL HAZARDS

Cover, collect, or incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

SAFETY RULES

IMPORTANT! Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is a danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP and get help right away.

CASORON GRANULES will selectively control annual grassy and certain perennial weeds around established ornamental trees and shrubs, fruit and nut trees, and berries, listed on this label. One application, made as directed, is generally enough to give season-long control.

CASORON GRANULES when applied to the soil as directed will give selective control of annual grassy and broadleaf weeds and certain perennial weeds such as:

- |              |                  |                  |
|--------------|------------------|------------------|
| Annual       | Fiddleneck       | Purslane         |
| Bluegrass    | Field Morning    | Quackgrass (from |
| Artemesia    | Glory            | seed)            |
| Bluegrass    | Florida Purslane | Ragweed          |
| Bull Thistle | (Pusley)         | Red Deadnettle   |

**BEST AVAILABLE COPY**

Camphorweed	Foxtail	Redroot Pigweed
Canada Thistle	Gisekia	Rosarypea
Carpetweed	Groundsel	Russian Knapweed
Chickweed	Henbit	Russian Thistle
Citron Melon	Horsetail	Shepherdspurse
Coffeeweed	Jerusalem oak	Smartweed
Crabgrass	Goosefoot	Spanishneedles
Cudweed	Knotweed	Spurge
Curly Dock	Lambsquarters	Teaweed
Dandelion	Milkweed Vine	Texas Panicum
Dogfennel	(Morrenia	(Hurrahgrass)
Evening	odorata)	Wild Carrot
Primrose	Natalgrass	Wild Mustard
False	Peppergrass	Wild Radish
Dandelion	Pineapple Weed	Yellow Woodsorrel
	Plantain	

CASORON GRANULES are to be applied only to established woody plants - those which have been planted for at least four months.

CASORON GRANULES can be used for weed control on the following Ornamental Woody Shrubs:

Amur Corktree	English Walnut	Mock Orange
Arborvitae	Euonymus	Mountain Ash
Ash	Filbert	Nandina
Azalea 1	Flowering Almond	Oak
Barberry	Flowering Quince	Osmanthus
Beauty Bush	Forsythia	(Holly-Olive)
Birch	Gardenia	Pachistima
Box Elder	Goldenraintree	Pecan
Boxwood	Hackberry	Photinia
Camellia	Heather	Pittosporum
Caragana	Holly 2	Poplar
Cleyera	Honeysuckle	Privet
Japonica	Ivy (Hedera Helix)	Pyracantha
Lotoneaster	Juniper	Rhododendron
Cottonwood	Pannikinnick	Rockrose
Crabapple	Laurel	Rose
Deutzia	Leucothoe	Russian Olive
Dogwood	Lilac	Spiraea
Eastern Red	Linden	Squaw-carpet
Cedar	Locust	Weigela
Elm	Magnolia	Willow
Elwood Cypress	Maple	Yew (Taxus)

1 Kurume, Mollis, Hardy Hybrid types and hardy native species only.  
 2 Except I. Crenata, I. Rotunda, I. Vomitoria.

CASORON GRANULES may be used for weed control on the following Edible Fruits & Nuts:

Apple	Filbert	Pear
Blackberry	Grape	Prune

**BEST AVAILABLE COPY**

Blueberry  
Cherry

Nectarine  
Peach

Plum  
Raspberry

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Before Application

1. Remove existing weeds.
2. Thoroughly work and smooth the area to be treated.
3. Carefully measure area to be treated and determine the number of square feet.
4. If area to be treated is large, it should be divided into smaller sections and each treated individually.

#### Application and Rate

1. Shake CASORON GRANULES uniformly on the soil at the rates specified.

FOR ANNUAL WEEDS: Apply in early spring before weeds emerge or after removing weeds.

FOR PERENNIAL WEEDS: For best control, apply after a killing frost from late fall to early winter.

#### 2. ORNAMENTAL WOODY SHRUBS:

Use at rate of 3/4 ounce (2 level tablespoons) for each 10 sq. ft. to be treated. 10 pounds treats 2,133 sq. ft.

FRUIT TREES, NUTS, BLUEBERRIES, GRAPES: Apply 2-3 level tablespoons per 10 sq. ft. Use the higher rate - 3 tablespoons per 10 sq. ft. if weed infestation is heavy or if most weeds are perennial. Wait 30 days after making application before harvesting stone fruits or nuts.

BLACKBERRIES, RASPBERRIES: Use 1 level tablespoons per 10 sq. ft. Use only on established plantings. Apply before new shoots emerge in spring.

1. Use the diagram below as a guide for uniform distribution of granules.

#### DIAGRAM

4. Rate higher than those indicated may cause injury.
5. More accurate and uniform application can be obtained by applying one-half of the required material over the entire treatment area and repeating the application with the remaining material.
6. CASORON GRANULES should not be used on sloping areas, because runoff may result in injury to lawn or other plants in adjacent areas.

#### After Application

1. Unless the soil is wet, sprinkle lightly to moisten the soil 1/2 to 1 inch deep. Do not flood, avoid puddling and washing.
2. Avoid disturbing soil in the treated area as long as weed control is being maintained.

**BEST AVAILABLE COPY**

NOTE: Use only on the plants listed. Do not apply to transplant or cutting beds or to newly set transplants, and do not seed or transplant into treated areas for one year after application. Do not use on extremely light sandy soils. Do not plant rotational crops, on which Dichlobenil is not registered, in treated soil within one (1) year after application.

**STORAGE AND DISPOSAL**

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse bag. Securely wrap bag in several layers of newspaper and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Filled by weight not by volume. Product may settle after packaging.

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON 97218  
EPA Reg. No. 801-530  
EPA Est. 802-OR-1  
29807