

PM 11:00 AM 9/1

LILLY/MILLER
Casoron Granules

Stops Weeds & Grasses From Growing Around Roses, Shrubs, Evergreens, Fruits & Berries

New refillable container

ACTIVE INGREDIENT:


Dichlobenil: 2,6-Dichlorobenzonitrile 2.0%
INERT INGREDIENTS 98.0%

Trademark and research discovery of Duphar, B.V., Amsterdam, Holland.

TREATS OVER 2,000 SQ. FT.

Keep out of reach of children
CAUTION
See back panel for additional cautions

NET WEIGHT 10 LBS.



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Do not allow contact with eyes or skin. In case of contact, flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Avoid contamination of feed and foodstuff.

ENVIRONMENTAL HAZARDS

Cover, collect, or incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

SAFETY RULES

IMPORTANT! Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is a danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP and get help right away.

CASORON GRANULES will selectively control annual grassy and certain perennial weeds around established ornamental trees and shrubs, fruit and nut trees, and berries, listed on this label. One application, made as directed, is generally enough to give season-long control.

CASORON GRANULES when applied to the soil as directed will give selective control of annual grassy and broadleaf weeds and certain perennial weeds such as:

- | | | |
|-----------|---------------|------------------|
| Annual | Fiddleneck | Purslane |
| Bluegrass | Field Morning | Quackgrass (from |

Artemesia	Glory	seed)
Bluegrass	Florida Purslane	Ragweed
Bull Thistle	(Pusley)	Red Deadnettle
Camphorweed	Foxtail	Redroot Pigweed
Canada Thistle	Gisekia	Rosarypea
Carpetweed	Groundsel	Russian Knapweed
Chickweed	Henbit	Russian Thistle
Citron Melon	Horsetail	Shepherdspurse
Coffeeweed	Jerusalem oak	Smartweed
Crabgrass	Goosefoot	Spanishneedles
Cudweed	Knotweed	Spurge
Curly Dock	Lambsquarters	Teaweed
Dandelion	Milkweed Vine	Texas Panicum
Dogfennel	(Morrenia	(Hurrahgrass)
Evening	odorata)	Wild Carrot
Primrose	Natalgrass	Wild Mustard
False	Peppergrass	Wild Radish
Dandelion	Pineapple Weed	Yellow Woodsorrel
	Plantain	

CASORON GRANULES are to be applied only to established woody plants - those which have been planted for at least four months.

CASORON GRANULES can be used for weed control on the following Ornamental Woody Shrubs:

Amur Corktree	English Walnut	Mock Orange
Arborvitae	Euonymous	Mountain Ash
Ash	Filbert	Nandina
Azalea 1	Flowering Almond	Oak
Barberry	Flowering Quince	Osmanthus
Beauty Bush	Forsythia	(Holly-Olive)
Birch	Gardenia	Pachistira
Box Elder	Goldenraintree	Pecan
Boxwood	Hackberry	Photinia
Camellia	Heather	Pittosporum
Caragana	Holly 2	Poplar
Cleyera	Honeysuckle	Privet
Japonica	Ivy (Hedera Helix)	Pyracantha
Cotoneaster	Juniper	Rhododendron
Cottonwood	Kinnikinnick	Rockrose
Crabapple	Laurel	Rose
Deutzia	Leucothoe	Russian Olive
Dogwood	Lilac	Spirea
Eastern Red	Linden	Squaw-carpenter
Cedar	Locust	Weigela
Elm	Magnolia	Willow
Elwood Cypress	Maple	Yew (Taxus)

1 Kurume, Mollis, Hardy Hybrid types and hardy native species only.
 2 Except I. Crenata, I. Rotunda, I. Vomitoria.

CASORON GRANULES may be used for weed control on the following Edible Fruits & Nuts:

Apple
Blackberry
Blueberry
Cherry

Filbert
Grape
Nectarine
Peach

Pear
Prune
Plum
Raspberry

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before Application

1. Remove existing weeds.
2. Thoroughly work and smooth the area to be treated.
3. Carefully measure area to be treated and determine the number of square feet.
4. If area to be treated is large, it should be divided into smaller sections and each treated individually.

Application and Rate

1. Shake CASORON GRANULES uniformly on the soil at the rates specified.

FOR ANNUAL WEEDS: Apply in early spring before weeds emerge or after removing weeds.

FOR PERENNIAL WEEDS: For best control, apply after a killing frost from late fall to early winter.

2. ORNAMENTAL WOODY SHRUBS:

Use at rate of 3/4 ounce (2 level tablespoons) for each 10 sq. ft. to be treated. 10 pounds treats 2,133 sq. ft.

FRUIT TREES, NUTS, BLUEBERRIES, GRAPES: Apply 2-3 level tablespoons per 10 sq. ft. Use the higher rate - 3 tablespoons per 10 sq. ft. if weed infestation is heavy or if most weeds are perennial. Wait 30 days after making application before harvesting stone fruits or nuts.

BLACKBERRIES, RASPBERRIES: Use 2 level tablespoons per 10 sq. ft. Use only on established plantings. Apply before new shoots emerge in spring.

3. Use the diagram below as a guide for uniform distribution of granules.

DIAGRAM

4. Rates higher than those indicated may cause injury.
5. More accurate and uniform application can be obtained by applying one-half of the required material over the entire treatment area and repeating the application with the remaining material.
6. CASORON GRANULES should not be used on sloping areas, because runoff may result in injury to lawn or other plants in adjacent areas.

After Application

1. Unless the soil is wet, sprinkle lightly to moisten the soil 1/4 to 1/2

- inch deep. Do not flood, avoid puddling and washing.
2. Avoid disturbing soil in the treated area as long as weed control is being maintained.

NOTE: Use only on the plants listed. Do not apply to transplant or cutting beds or to newly set transplants, and do not seed or transplant into treated areas for one year after application. Do not use on extremely light sandy soils. Do not plant rotational crops, on which Dichlobenil is not registered, in treated soil within one (1) year after application.

REFILLING INSTRUCTIONS:

- 1/ Be sure that the top applicator ring is in the closed position.
- 2/ Set the canister upside down on a firm surface.
- 3/ Remove bottom "plug" with a coin or screwdriver.
- 4/ Carefully refill canister with Lilly/Miller Casoron Granules.
- 5/ Press "plug" firmly back into position.
- 6/ Return the canister to the upright position.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container (bag), except as directed for refilling. Securely wrap container (bag) in several layers of newspaper and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Filled by weight not by volume. Product may settle after packaging.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

EPA Reg. No. 802-536

EPA Est. 802-OR-1 (P)

EPA Est. 407-IA-1 (S)

EPA Est. 279-CA-1 (F)

Superscript used is first letter of lot number
29807