



LILLY/MILLER  
TOMATO & VEGETABLE GARDEN SPRAY

ILLUSTRATION

802-540

ACTIVE INGREDIENTS:

Pyrethrins.....	0.050%
*Piperonyl Butoxide, Technical.....	0.500%
Petroleum Distillate.....	8.075%
INERT INGREDIENTS:.....	91.375%
	<u>100.000%</u>

\*Equivalent to 0.40% (Butylcarbityl) (6-propylpiperonyl) ether and 0.10% related compounds

PYRENONE- Registered trademark of Fairfield American Corporation

An Effective Fast Acting Insecticide For Use On Vegetables.

Accepted for Use to Day of Harvest on Edible Plants

<p>KEEP OUT OF REACH OF CHILDREN</p> <p>CAUTION</p> <p>See side panels for additional precautionary labeling.</p>
---

10 pt.

12 pt.

NET WT. 14 OZ.

.....  
.....  
.....  
.....

Left Side Panel:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Keep out of reach of children. Avoid breathing of spray or contact with eyes or skin. Wash thoroughly after use. Do not contaminate feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not use near fishbowls or aquaria. Keep out of lakes, streams or ponds.

PHYSICAL HAZARDS

Contents under pressure. Do not puncture. Do not store near heat or open flame. Do not store below freezing temperatures. Exposure to temperatures above 130° F may cause bursting. Do not throw container into fire or incinerator.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Wrap container and dispose of in trash. Do not incinerate.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

LC-1Q

THE CHAS. H. LILLY CO.  
Portland, Oregon 97218

EPA Reg. No. 802-

EPA Est. No. 10900-CA-1 (S) EPA Est. No. 11525-CA-1(F)

Superscript used is first letter of lot  
number

Right Side of Panel:

#### USE DIRECTIONS

SHAKE CAN WELL BEFORE USING.

This insecticide may be used outdoors in gardens throughout the growing season or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

For best results, use throughout the growing season to prevent buildups of insect infestations. Spray plants from a distance of 1½ to 2 feet using short bursts of 1 second. To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when the air is still, treating all foliage with particular attention to the undersides of leaves. Contact insects directly whenever possible. Repeat treatment every 7-10 days or as reinfestation occurs.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatments as needed to control infestations. May be applied on day of harvest on the following crops to control the listed insects:

ASPARAGUS:	Asparagus Beetle
BEANS:	Aphids, Leafhoppers, White fly, Mexican Bean Beetles, 12-Spotted Cucumber Beetles.
BROCCOLI, CABBAGE, BRUSSEL SPROUTS, CAULIFLOWER:	Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug.
CELERY:	Green Peach Aphid, Cabbage Looper, Leaf-tier.
CRANBERRIES:	Fireworms, Leafhoppers.
CUCUMBERS, MELONS, SQUASH:	Flea Beetles, Green Peach Aphids.
EGGPLANTS:	Blister Beetles, Colorado Potato Beetles, Flea Beetles, Green Peach Aphids.
LETTUCE:	Cabbage Looper, Green Peach Aphids, Diamondback Moth Larvae, Imported Cabbageworms.
MUSTARD GREENS, KALE, COLLARDS, TURNIPS:	Imported Cabbageworms, Diamondback Caterpillars, Aphids, Cabbage Looper.
PEPPERS:	Green Peach Aphids.
POTATOES:	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers.
RADISHES:	Aphids, Flea Beetles.
SPINACH:	Aphids, Cabbage Looper, Webworms.
SWEET CORN:	Japanese Beetles.
TOMATOES:	Flea Beetles, Colorado Potato Beetles, Stink Bugs, Green Peach Aphids.