


Front Panel

LILLY/MILLER
DEVIRINOL^R 2-G
Pre-emergent Herbicide

Illustration



A Pre-emergence Herbicide for Weed Control in Home Flower Gardens and Ornamental Plantings. Use around Deciduous and Evergreen Trees and Shrubs, Ground Covers, Perennial Flowers, Fruit and Nut Crops and Container Grown Plants.

ACTIVE INGREDIENT

N,N-Diethyl-2-(1-naphthalenyloxy)-propionamide 2.0%

INERT INGREDIENTS

98.0%

100.0%

^RTrademark of Stauffer Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WT. 5 LB., 20 LB.

Right Side of Panel

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHAT THIS PRODUCT DOES

Lilly/Miller DEVRINOL 2-G controls certain annual grassy and broadleaf weeds as they germinate. It does not control established weeds.

See list of weeds controlled on left side panel.

WHERE TO USE THIS PRODUCT

Lilly/Miller DEVRINOL 2-G can be applied over small ornamental plants or around larger plants in rose gardens, container plantings, ornamental landscapes and ground covers including dichondra.

Use Lilly/Miller DEVRINOL 2-G for weed control around the following plants

DECIDUOUS AND EVERGREEN TREES

Acer	Fir	Picea
Ash	Gleditsia	Pinus
Black Walnut	Hemlock	Pittosporum
Cedar	Japanese Larch	Podocarpus
Citrus	Laburnum	Populus
Crateagus	Malus	Prunus
Cypress	Olive (Olea)	Pears
Douglas Fir	Palm (Washingtonia	Pistachios
Eucalyptus	rebusta)	
Ficus		

DECIDUOUS AND EVERGREEN SHRUBS

Abelia	Hypericum	Pittosporum
Asparagus (Sprengen)	Ilex	Podocarpus
Azalea	Juniperus	Pyracantha
Bottlebrush	Ligustrum	Rhododendron
Buxus	Nandina	Rosa
Euonymus	Natal Plum	Star Jasmine
Hibiscus	Oleander	Taxus
	Photinia	

GROUND COVERS

Ajuga	Gazania	Liriope
Carpobrotus	Hedera	Osteosperma
Delosperma	Hypericum	Pachysandra
Dichondra	Lantana	Vinca
Erysimum		

Right Side of Panel - Cont'd

CONTAINER STOCK

Abelia	Fir	Photinia
Agapanthus	Hedera	Pinus Spp.
Asparagus	Hibiscus	Pittosporum
Azalea	Ilex	Podocarpus
Bottlebrush	Juniperus	Pyracantha
Buxus	Liriodendron	Rhododendron
Cedar	Ligustrum	Rosa
Cypress	Nandina	Star Jasmine
Douglas Fir	Pachysandra	Thuja
Eucalyptus	Palm (Washingtonia robusta)	Vinca
Euonymus		

PERENNIAL FLOWERS

African Daisy	Bird of Paradise	Chrysanthemum
---------------	------------------	---------------

FRUIT & NUTS

Almond	Lemon	Pistachio
Apple	Nectarine	Plum
Apricot	Orange	Prune
Cherry	Peach	Tangerine
Filbert	Pear	Walnut
Grape	Pecan	

Strawberry (Application should be delayed until the desired number of daughter plants have become established. Do not apply from bloom to harvest.)

WHEN TO USE THIS PRODUCT

Lilly/Miller DEVRINO, 2-G should be used before weeds germinate. It will not control established weeds. Application can be made anytime of the year to weed-free soil. Fall and winter applications will provide excellent weed control the following spring and summer. In containers, apply after setting and watering the plants.

HOW TO USE THIS PRODUCT

Apply this product as specified on this label. Lilly/Miller DEVRINOL 2-G should be spread evenly on the soil surface using shaker can or granular herbicide applicator. Do not apply when weather conditions favor drift from areas treated.

When weed infestation is heavy or when broadleaves are the primary weeds, use one pound on 145 square feet (6-9 lbs/1,000 sq. ft.) For annual grassy weeds or when weed infestation is light, use one pound on 218 square feet (4-6 lbs/1,000 sq. ft.)

Lilly/Miller DEVRINOL 2-G should be incorporated evenly into the soil for optimum results. Use a hand cultivator to a depth of 1-2 inches or water thoroughly (1-2 inches) within 2-3 days.

Left Side Panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: MAY BE HARMFUL IF SWALLOWED. Do not breathe dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after use. Avoid Contamination of Food and Feed.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, flush with plenty of water and get medical attention.

IF ON SKIN, flush with plenty of water for at least 15 minutes and get medical attention if irritation persists. Remove contaminated clothing and shoes. Wash clothing thoroughly before reuse.

IF SWALLOWED, immediately dilute the swallowed material by giving large quantities of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Continue fluid administration until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. Seek medical attention if respiratory irritation occurs.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

STORAGE: Keep away from food, feedstuffs, and domestic water supplies.

DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Wrap empty container and put in trash collection.

Left Side of Panel - Cont'd

WEEDS CONTROLLED BY DEVRINOL 2-G

ANNUAL GRASSES:

Barnyardgrass, Junglerice

Wild Barley

Annual Bluegrass

Bromes

Cheatgrass

Downy Brome

Ripgut Brome

Soft Chess

Large (Hairy) Crabgrass

Smooth Crabgrass

Southwestern Cupgrass

Foxtails

Bristly Foxtail

Giant Foxtail

Green Foxtail

Yellow Foxtail

Goosegrass

Seeding Johnsongrass

Wild Oats

Panicums

Guineagrass

Fall Panicum

Texas Panicum

Witchgrass

Italian Ryegrass

Sandbur

Red Sprangletop

Stinkgrass

ANNUAL BROADLEAF WEEDS:

Carpetweed

Common Chickweed

Coast Fiddleneck

Filaree

Groundsel

Prostrate Knotweed

Lambsquarters

Prickly Lettuce

Little Mallow

Redroot Pigweed

Pineappleweed

Common Purslane

Horse Purslane

Common Ragweed

Annual Sowthistle

From Seed

Suppression