

p m 25

802-545

1 of 4

LILLY/MILLER
DEVIRINOL® 2-G
Pre-emergent Herbicide
Illustration

ACCEPTED

MAR - 5 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 802-545

Controls weeds as they sprout in Flower Gardens,
Ornamental Plantings and Home Orchards

ACTIVE INGREDIENT:

N,N-Diethyl-2-(1-naphthalenyloxy)-propionamide	2.0%
INERT INGREDIENTS	98.0%
TOTAL	<u>100.0%</u>

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Wt. 1 1/4 lb., 5 lbs.

PRECAUTIONARY STATEMENTS

2 of 4

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: MAY BE HARMFUL IF SWALLOWED. Do not breathe dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after use. Avoid Contamination of Food and Feed.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, flush with plenty of water and get medical attention.

IF ON SKIN' flush with plenty of water for at least 15 minutes and get medical attention if irritation persists. Remove contaminated clothing and shoes. Wash clothing thoroughly before reuse.

IF SWALLOWED, immediately dilute the swallowed material by giving large quantities of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue.

Continue fluid administration until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. Seek medical attention if respiratory irritation occurs.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Lilly/Miller DEVRINOL 2G provides selective preemergent control of annual grassy and broadleaf weeds around trees, shrubs, ground cover, listed flowers and around fruit and nut trees. Use Lilly/Miller Devrinol 2G to control annual weeds before they germinate. For best results, apply once in fall or early winter to insure good weed control during the following spring and summer.

Make applications around the following ornamental and edible plants:

ORNAMENTAL TREES & SHRUBS

Abelia	Fig	Oleander
Ash	Fir	Olive (Olea sp.)
Asparagus (Spenger)	Firethorn	Palm (Washingtonia robusta)
Aspen	Hawthorn	Pear
Azalea	Hemlock	Photinia
Black Walnut	Hibiscus	Pine
Bottlebrush	Holly	Pistachio
Boxwood	Honey Locust	Pittosporum
Cedar	Hypericum	Podocarpus
Citrus	Japanese Larch	Poplar
Cottonwood	Juniper	Privet
Crabapple	Laburnum	Rhododendron
Cypress	Maple	Rose
Douglas Fir	Nandina	Spruce
Eucalyptus	Natal Plum	Star Jasmine
Euonymous		Yew

GROUND COVERS

3 of 4

Ajuga
African Daisy
Delosperma
Dichondra
Erysimum

Gazania
Hypericum
Ice Plant
Ivy
Lantana

Liriope
Myrtle
Pachysandra
Periwinkle

PERENNIAL FLOWERS

African Daisy

Bird of Paradise

Chrysanthemum

EDIBLE FRUIT & NUTS

Almond
Apple
Apricot
Cherry
Filbert
Grape

Lemon
Nectarine
Orange
Peach
Pear
Pecan

Pistachio
Plum
Prune
Tangerine
Walnut

Strawberry (Application should be delayed until the desired number of daughter plants have become established. Do not apply from bloom to harvest.)

WEEDS CONTROLLED BY DEVRINOL 2-G

ANNUAL GRASSES:

Barnyardgrass, Junglerice
Wild Barley
Annual Bluegrass
Bromes
 Cheatgrass
 Downy Brome
 Ripgut Brome
 Soft Chess
Large (Hairy) Crabgrass
Smooth Crabgrass
Southwestern Cupgrass
Foxtails
 Bristly Foxtail
 Giant Foxtail
 Green Foxtail
 Yellow Foxtail
Goosegrass
Seeding Johnsongrass
Wild Oats
Panicums
 Guineagrass
 Fall Panicum
 Witchgrass
Italian Ryegrass
Sandbur
Red Sprangletop
Stinkgrass

ANNUAL BROADLEAF WEEDS:

Carpetweed
Common Chickweed
Coast Fiddleneck
Filaree
Groundsel
Prostrate Knotweed
Lambsquarters
Prickly Lettuce
Little Mallow *
Redroot Pigweed
Pineappleweed
Common Purslane
Horse Purslane
Common Ragweed **
Annual Sowthistle
 *From Seed
 **Suppression

4.84

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Before Application

1. Lilly/Miller Devrinol®2G controls annual grassy and broadleaf weeds as they germinate. Because it does not control established weeds, turn up 1-2 inches of soil in the area to be treated and remove the weeds.
2. Smooth out the area and determine the number of square feet to be treated.
3. If the area is large, divide it into smaller areas and treat each area individually.

Application and Rate

1. Shake Lilly/Miller Devrinol®2G uniformly over the soil using shaker canister or granular herbicide applicator. Do not apply when weather conditions favor drift from area treated.
2. If area had heavy weed infestation prior to treatment, or if most weeds were broadleaf, use 1 lb. over 145 sq. ft. (7 lbs. per 1000 sq. ft.)
3. If area had light weed infestation or if most weeds were annual grassy weeds, use 1 lb. over 218 sq. ft. (4.5 lbs. per 1000 sq. ft.)

After Application

For best results, mix Devrinol®2G into the top 1-2 inches of soil, or water in thoroughly within 2-3 days.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals.

Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

•Trademark of Stauffer Chemical Company

The Chas. H. Lilly Co.
Portland, Oregon 97218

EPA Reg. No. 802-545

EPA Est. 802-CR-1