ULTRA GREEN

WEED & HIED

Illustration

Performance formula controls weeds and provides a thick, healthy fawn in one application

21-3-5

CONTAINS TRIMEC HERBICIDE

COVERAGE Up to 4,000 sq. ft.

ACCEPTED JUNI 1 1534 ACTIVE INGREDIENTS:
Dimethylamine Salt of 2,4 Dichlorophenoxyacetic acid

0.44

Dimethylamine Salt of 2-(2-methyl-4 chlorophenoxy) propionic acid. 0.28

Dimethylamine Salt of Dicamba (3.6-dichloro O anisic acid)*** 0.14

INERT INGREDIENTS: 99.14% ___

sale.

0.236%

BEST DOCUMENT AVAILABLE

•2,4 Dichlorophenoxyacetic acid isomer specific by AOAC Method 6,002.5

6.D02 **5.** 0.3

**2 (2 methyl 4 chlo; opherioxy) = propionic acid

***3,6 Dichloro o anisic acid 0.119%

TRIMEC Reg. IM of PBI/Gordon Corp., U.S. Pat. No. 3,784,186

NET WT. 20 lbs.

Keep out of reach of children

CAUTION

See back panel for edditional processionary statements.

800 541

LILLY/MILLER

ULTRA GREEN WEED & FEED Illustration

The performance story.

Ultra Green Weed & Feed is a specially selected combination of plant foods and herbicide formulated to provide satisfactory results with one application.

Water Soluble Nitrogen goes to work immediately after watering. Nitrogen works as a protein builder and helps grass to grow healthy and strong with a rich green > color.

Time release Nitrogen works slowly and continuously over time as it is released by bacterial action in the soil. It huilds protein and color through steady feeding.

Phosphoric acid is an important nutrient in early growth and root formation. It helps young plants grow strong.

Potash encourages a strong root system and helps build resistance to disease,

Sulphur is an essential element for protein movement. It helps nitrogen promote strong, sturdy growth.

Iron is important for chlorophyll production. It helps turn your lawn that incredibly deep shade of green.

Zinc helps regulate plant growth. Zinc works with nitrogen to build plant protein.

TRIMEC "Herbicide is a patented combination of three proven weed killers that, together, control a wide range of broadle af lawn weeds. Including databelions, plantain, chickwood, clover, Japanese clover, herbit, buttercop, spurge, lawn daisy, puncture vine and others.

PAST THE UNLINE AVAILABLE

For Best Results:

- 1. Do not mow lawn for two days prior to application.
- 2. Do not apply to new bans under 3 months old.
- 3. Apply to moist soil and foliage using an accurate spreader and the appropriate dial setting from chart below. Do not water for at least 24 hours after application.
- A second application can be applied after one month to control any particularly hard to kill weeds.

HOW TO SPREAD LAWN FERTILIZER (Illustrations)

- 1. Begin spreading by laying down two strips of lawn food at each end of your lawn. I ill in remaining area as shown here.
- 2. Make turns with the spreader CLOSED to avoid uneven coverage.
- Overlap previous wheel track slightly, as shown above to avoid gaps in coverage.
- 4. Close the spreader when you stop walking to avoid uneven coverage. Stopping and starting with spreader box open will waste your fawn food.

SPREADER SETTINGS

	Filly/Miller Spreader	Contral Spreader	Turf King	Scotts Model 35 & 75	
Lilly/Miller Ultra Green Weed & Feed	81/2	1½	8	81/2	

NOTICE

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking queed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co., nor the retail dealer can generable exact coverage as indicated by the recommended dial settings.

BEST DOCUMENT AVAILAE .

GUARANTEED FERTILIZER ANALYSIS

Fotal Nitrogen (N)	O'
2.0% Ammoniacal Nitrogen	
(From Ammonium Phosphate)	
15.5% Urea Nitrogen (from Urea and Sulphur-Coated Urea)	
3.5% Coated Slow Release Urea Nitrogen	
(from Sulphur Coated Urea)	
Available Phosphoric Acid (P ₂ O ₅)	%
Soluble Potash (K ₂ O)	%
Iron (Fe)	%

C. J. C. 188 .

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domostic Animals

caurion: Harrful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact irrediately flush eyes 5217.339 or skin with plenty of water. Get redical attention if irritation persists.

STATEMENT OF PRACTICAL TREATMENT

In case of contact with eyes, flush with water for 15 minutes. Get redical attention. In case of contact with skin, wash with plenty of soap and water.

ENVIRORMENTAL HAJARDS

Do not apply directly to water. Do not contaminate domestic or irritation waters. Do not apply when weather conditions favor frift from target area. Do not contaminate water by cleaning of equapment or disposal of waste.

- T DOCUMIENT AVAILABLE

3. No Col 1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection. Keep pesticide in original container. Keep container closed when not in use. Do not store in contact with seeds or other pesticides. Do not store near feed or food stuffs.

USE PRECAUTIONS

2,4 D, MCPP and Banvel are injurious to most plants other than grasses, so this material should be applied carefully on a day with low wind velocity to avoid drift onto valuable plants. Do not use on St. Augustine grass, Dichondra, carpetgrass or clover lawns. Can cause possible injury to bentgrass. Do not apply within the drip line of desirable trees and shrubs. Rinse spreader with clear water after use.

NOTICE: Follow directions carefully. Firning and method of application, weather conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LC-1S

The Chas. H. 1 illy Co. Portlad, Oregon 97218

LPA Reg. No. 802-547,

1PA Est. No. 802 OR-1

WEST DOCUMENT AVERAGES