

The
Chas. H.
LILLY Co.



PM 23

802-547

144

7737 N.E. KILLINGSWORTH • PORTLAND, OREGON 97218 • (503) 256-4600
• FAX (503) 255-7552

February 27, 1993

I, being an authorized representative of The Chas. H. LILLY Co., certify that all containers of Lilly/Miller ULTRAGREEN Weed & Feed, EPA Reg. No. 802-547 produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Molly Klettke
Molly Klettke, Registration Specialist

299

LILLY/MILLER ULTRAGREEN WEED & FEED

Kills Dandelions, Chickweed, Knotweed, Plantains & Other Broadleaf Weeds
In Lawns

Greens-up your lawn quickly

Covers up to 4,000 square feet

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid*	0.44%
INERT INGREDIENTS	99.56%

*2,4-Dichlorophenoxyacetic acid equivalent 0.370%. Isomer specific by
AOAC Method

CAUTION. Keep out of reach of children. See back panel for additional
precautionary statements.

Net Weight 20 lbs.

) PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. When using this product, wear long
pants, long-sleeved shirt, socks and shoes. After using this product,
remove clothing and launder separately before reuse, and promptly and
thoroughly wash hands and exposed skin with soap and water. Avoid
contact with skin, eyes or clothing. In case of contact immediately
flush eyes or skin with plenty of water. Get medical attention if
irritation persists. Do not allow people (other than applicator) or
pets on treatment area during application. Do not reenter or permit
others to reenter treated areas until granules are washed into the soil
and grass has dried.

STATEMENT OF PRACTICAL TREATMENT

In case of contact with eyes: Flush with water for 15 minutes. Get
medical attention. In case of contact with skin: Wash with plenty of
soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may
adversely affect aquatic invertebrates and nontarget plants. Do not
apply directly to water or wetlands (swamps, bogs, marshes, and
potholes). Do not contaminate water when disposing of equipment
washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling. The maximum number of broadcast
applications per treatment site is 2 per year.

Use Precautions

2,4-D is injurious to most plants other than grasses, so this material
should be applied carefully on a day with low wind velocity to avoid
drift onto valuable plants. Applications during hot weather (over 80
degrees F) are apt to cause some temporary damage to turf. Do not use
on St. Augustine grass, Dichondra, carpetgrass or clover lawns. Can
cause possible injury to bentgrass. Do not apply within the drip line

of desirable trees and shrubs. Rinse spreader with clear water after use.

STORAGE AND DISPOSAL

Store this product in its original container out of reach of children and domestic animals. Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

Do not mow lawn for two days prior to application. Do not apply to new lawns under three months old. Apply to moist soil and foliage using an accurate spreader and the appropriate dial setting from the chart below. Do not water for at least 24 hours after application. A second application can be applied after one month to control any particularly hard-to-control weeds.

(Captions for illustrations)

) Lawns must be thoroughly watered the day before application. Then water again lightly, just prior to application. Begin spreading by laying down two strips of lawn food at each end of lawn. Fill in remaining area as shown here. Close the spreader when you stop walking or make turns to avoid uneven coverage. Stopping and starting with the spreader box open will waste lawn food. Overlap previous wheel track slightly, as shown, to avoid gaps in coverage. Wait 24 hours for herbicide to work. Then, water in to start the greening process.

Spreader Settings

Application Rate 20 lbs. per 4,000 sq. ft.

Lilly/Miller Spreader	6
Central Spreader	5
Scotts Model 35, 75 & PF-1	6
Ortho 3000	4 1/2
Other models	To be determined

) Notice: Spreader settings given have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

Notice: Follow directions carefully. Timing and method of application, weather conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N) 20%

17.6% Ammoniacal Nitrogen

2.4% Urea Nitrogen

Available Phosphoric Acid (P2O5) 2%

Soluble Potash (K2O) 5%

Sulfur (S) 19%

Derived from Ammonium Sulfate, Ammonium Phosphate, Urea, and Muriate of

4 of 4

Potash.

Filled by weight not by volume. Product may settle after packaging.

The Lilly/Miller Performance Guarantee

If you are not satisfied with the performance of this product, simply send your name, address, and proof of purchase to The Chas. H. LILLY Co., 7737 NE Killingsworth, Portland, OR 97218, and we will be happy to refund your money.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Reg. No. 802-547
EPA Est. 802-WA-1

