LILLY/MILLER DEVRINOL*5-G

L

A PRE-EMERGENCE, SELECTIVE HERBICIDE FOR WEED CONTROL IN CONTAINER GROWN ORNAMENTALS, FIELD-GROWN NURSERY STOCK, LINER STOCK AND GROUND COVERS.

	ACTIVE INGREDIENT: N,N-Diethyl-2-(1 naphthalenyloxy)-propionamide		
	100.0%		
T	Contains 5 pounds Active Ingredient per 100 pounds.		
	[®] Trademark of Stauffer Chemical Company		
	KEEP OUT OF REACH OF CHILDREN		
	CAUTION		
	PRECAUTIONARY STATEMENTS		
	HAZARDS TO HUMANS AND DOMESTIC ANIMALS		
	CAUTION: May be harmful if swallowed. Do not breathe dust. Avoid contact with		
	skin, eyes and clothing. Wash thoroughly with soap and water after use. Avoid contamination of food and feed. STATEMENT OF PRACTICAL TREATMENT		
	In case of contact, immediately flush eyes or skin with plenty of water for at		
	least fifteen minutes while removing contaminated clothing and shoes. Call a physician. Wash clothing before reuse.		
	If swallowed, immediately dilute the swallowed material by giving large		
	quantities of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Continue fluid administration until vomitus is		
	clear. Never give anything by mouth to an unconscious person.		
	If inhaled, remove to fresh air. Seek medical attention if respiratory irritation		
	occurs.		
	ENVIRONMENTAL HAZARDS		
	Do not apply directly to lakes, streams, and/or ponds.		
	Do not contaminate water by cleaning of equipment or diposal of wastes.		
	Apply this product as specified on this label.		
	Do not apply when weather conditions favor drift from areas treated.		
	Do not contaminate irrigation water or water used for domestic purposes.		

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

••• •••

-- - -- - -

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep container closed when not in use. Store in a cool dry place. Do not store near seeds, fertilizer or foodstuffs.

PESTICIDE DISPOSAL: Wash application equipment after use. Pesticide or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act. CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

DIRECTIONS FOR USE

1. DEVRINOL 5-G does not contol established weeds. It must be applied to freshly weeded soil before weeds germinate or during fall and winter months.

 DEVRINOL 5-G can be applied to newly planted container stock after soil has settled from first watering, field grown nursery stock, dichondra and to established plants.
DEVRINOL 5-G needs mechanical incorporation, (such as,power tiller), adequate irrigation or natural moisture (rainfall or snow) (1-2 inches within 2-3 days) for optimum results.

DEVRINOL 5-G is recommended on several different container potting soil 4. mixtures. These mixtures include those containing 30 percent soil and/or sand and mixtures containing no soil (Example:perlite, peat, bark, sawdust and sand) WEEDS CONTROLLED BY DEVRINOL 5-G ANNUAL GRASSES Barnyardgrass, Jungierice (Echinochioa spp) Wild Barley (Hordeum leporinum) Annual Bluegrass (Poa annua) Bromes (Bromus) Cheatgrass (B. secalinus) Downy Brome (B-tectorum) Ripgut Brome (B rigidus) Soft Chess (B. mollis) Large (Hairy) Crabtrass (Digitaria sanguinalis) Smooth Crabgrass (Digitaria ischaemum) Southwestern Cupgrass (Eriochloa gracilis) Foxtails (Setaria) Bristly Foxtail (S.verticilla) Giant Foxtail (S.faberi) Green Foxtail (Syindis) Yellow Foxtail (S lutescens) Goosegrass (Eleusine indica) Seedling Johnsongrass (Sorghum halepese) Wild Oats (Avena fatua) Panicums (Panicum) Guineagrass (P maximum) Fall Panicum (P dichotomiflorum) Texas Panicum (Pitexanum) Witchgrass (P capillare) Italian Ryegrass (Lolium multiflorum) Sandbur (Cenchrus spp.) Red Sprangletop (Leptochloa filiformis) Stinkgrass (Eragrostis cilianensis)

 ANNUAL BROADLEAF WEEDS Carpetweed (Mollugo verticillata) Common Chickweed (Stellaria media) Coast Fiddleneck (Amsinckia intermedia) Filaree (Erodium spp) Groundsel (Senecio spp) Prostrate Knotweed (Polygonum aviculare) Lambsquarters (Chenopodium album) Prickly Lettuce (Lactuca serriola) Little Mallow* (Malva parviflora) Redroot Pigweed (Amaranthus retroflexus) Pineappleweed (Matricaria matricarioides) Common Pursiane (Portulaca oleracea) Horse Purslane (Trianthema portullacastrum) Common Ragweed ** (Ambrosia artemisiifolia) Annual Sowthistle (Sonchus oleraceus) *From seed **Suppression

RECOMMENDATIONS

NURSERY STOCK—FIELD GROWN. Consult list of plant species on which DEVRINOL 5-G can be used. Apply DEVRINOL 5-G as a broadcast treatment over young nursery stock or as a directed application to larger stock either as a band or broadcast treatment. Application can be made anytime of the year to weed free soil, hoever, since DEVRINOL 5-G requires 1-2 inches of rainfall or irrigation soon after application for adequate weed control, application should be timed to take advantage of rainfall if irrigation is not available. Fall and winter applications will provide excellent weed control the following spring and summer.

A thorough incorporation of DEVRINOL 5-G into the soil to a depth of 1-2 inches will provide excellent weed control with less rainfall required.

One application of DEVRINOL 5-G should provide adequate weed control for a growing season in temperate areas.

CONTAINER GROWN ORNAMENTALS. Consult list of plant species on which DEVRINOL 5-G can be used. Apply DEVRINOL 5-G as a broad cast application over the containers.

Normal irrigation or rainfall (if a fall or winter application is made in cold areas) should provide sufficient water for satisfactory weed control.

HOME GROUNDS, HIGHWAY AND INDUSTRIAL ORNAMENTAL PLANTINGS. Use an overall (broadcast) treatment over young ornamentals or as a directed application where larger plants are growing. Irrigation is not available or if application is to be made in lower summer rainfall areas.

GROUND COVERS. Apply DEVRINOL 5-G as a broadcast treatment over the ground covers. Irrigate immediately with a minimum of 1 inch of water or apply when rainfall is expected.

DICHONDRA. Apply DEVRINOL 5-G st the time of seeding or to established dichondra including dichondra sod. Irrigate with a minimum of 1 inch of water.

APPLICATION RATES

ANNUAL WEEDS. For annual grasses and broadleaf weeds apply 80 to 120 lbs. of DEVRINOL 5 G per broadcast acre (1.8 to 2.8 lbs. per 1,000 sq. ft.)

Use the higher rate when weed populations are heavy or when broadleaves are the primary weeds.

USE DEVRINOL 5-G ON THESE ORNAMENTALS AND NURSERY STOCK AND CONTAINER STOCK

RIELD CROWN				
FIELD GROWN DECIDUOUS AND EVERGR	FEN TREES			
	Crataegus	Fir		
Acer	Cypress	Gleditsia		
Ash Diada Mala M	Douglas Fir	Hemlock		
Black Walnut	Eucalyptus	Japanese Larch		
Cedar	Ficus	Laburnum		
Cicrus	r içus			
Malus	Picea	Populus		
Olive (Olea)	Pinus	Prunus		
Palm	Pittosporum	Pears		
(Washington robusta)	Podocarpus	Pistachios		
(Washington / Obester	· • • • • • •			
DECIDUOUS AND EVERGREEN SHRUBS				
Abelia	Bottlebrush	Hypericum		
Asparagus	Buxus	llex		
(Sprenger)	Euonymus	Juniperus		
Azalea	Hibiscus	Ligustrum		
Nandina	Pittosporum	Rosa		
Natal Plum	Podocarpus	Star Jasmine		
Oleander	Pyracantha	Taxus		
Photinia	Rhododendron			
FRUIT AND NUTS				
Almond	Cherry	Lemon		
Apple	Filbert	Nectarine		
Apricot	Grape	Orange		
	Distochio	Tangerine		
Peach	Pistachio	Walnut		
Pear	Plum	4491191		
Pecan	Pecan Prune Prune			
Strawberry (Application should be delayed until the desired number of daughter plants have become established. Do not apply from bloom to harvest)				
become established. Do not	apply nom bleom to narvesty			
GROUND COVERS				
Ajuga	Dichondra	Gazania		
Carpobrofus	Erysimum	Hedera		
Delosperma				
Delosperina				
Hypericum	Liriope	Pachysandra		
Lantana	Osteosperma	Vinca		
CONTAINER STOCK				
Abelia	Cedar	Hedera		
Agapanthus	Cypress	Hibiscus		
Asparagus	Douglas Fir	llex		
Azalea	Eucalyptus	Juniperus		
Destabaseb	Fuonauus	Liriope		

Euonymus

Fir

Bottlebrush

Buxus

•

4

•

Liriope

CONTAINER STOCK cont.

Ligustrum Nandina Pachysandra Palm (Washington robusta)

PERENNIAL FLOWER

Bird of Paradise

Photinia Pinus Spp Pittosporum Podocarpus Pyracantha Rhododendron Rosa Star Jasmine Thuja Vinca

NOTICE. Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

LC-1R

...

EPA Est. No. 802-OR-1

The Chas. H. Lilly Co. Portland, Oregon 97218

EPA Reg. No. 802-

NET WT. 50 lbs.

•