


LILLY/MILLER
1½ CHLORBAN INSECT GRANULES

Alternate Names: Lilly/Miller Chlorban Lawn Insect Granules
Lilly/Miller Chlorban Insect Granules Contains
DURSBAN* Insecticide
Contains DURSBAN* Insecticide

LAWN INSECT CONTROL

- EUROPEAN CRANE FLY
- BILLBUGS
- ANTS
- CHINCH BUGS
- FLEAS
- GRASSHOPPERS



ACTIVE INGREDIENT:

Chlorpyrifos (0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl)
phosphorothioate) 1.0%
INERT INGREDIENTS 99.0%
Total100.0%

Keep out of reach of children

CAUTION

See back panel for Statement of Practical Treatment and additional
Precautionary Statements.

NET WEIGHT 10 LBS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.


FOR LAWNS AND OTHER ORNAMENTAL TURF AREAS

- | | |
|--------------|--------------------|
| To Control: | Billbugs |
| Ants | Brown Dog Ticks |
| Armyworms | Crickets |
| Chinch Bugs | Clover Mites |
| Chiggers | Earwigs |
| Cutworms | Fleas |
| Sod Webworms | Grasshoppers |
| (Lawn Moths) | Sowbugs (Pillbugs) |

Apply 1½ CHLORBAN INSECT GRANULES at the rate of 2 1/2 lbs. per 1,000
sq. ft. (1/2 lb. per 200 sq. ft.) of area. Use any hand operated
gravity spreader designed for application of granular materials. See
Spreader Settings.

Treat when insects first appear. Immediately after application,
sprinkle the turf with 6 gallons of water per 200 sq. ft. so as to wash
the granules into the zone of insect infestation.

WHITE GRUB CONTROL (Larvae of European Chafer and Japanese Beetle)



Apply 1½ CHLORBAN INSECT GRANULES at the rate of 5-10 lbs. per 1,000 square feet (1-2 lbs. per 200 square feet) of area.

See Spreader Settings. Apply the granules uniformly over the area treated, followed immediately by a thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service for proper time to treat in your area.

**EUROPEAN CRANE FLY LARVAE CONTROL
(Leather Jackets)**

Apply 3 1/2 lbs. 1½ CHLORBAN INSECT GRANULES to 1,000 square feet using any hand operated gravity spreader designed for application of granular materials. See Spreader Settings. Immediately after application, sprinkle the turf with 30 gallons of water per 1,000 square feet. For timing of applications you should consult your local Agricultural Extension Service.

HYPERODES (Turfgrass) WEEVIL CONTROL

Apply 1½ CHLORBAN INSECT GRANULES at the rate of 5 lbs. per 1,000 square feet (1 lb. per 200 square feet) of area.

See Spreader Settings. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service Specialist for best time to treat in your area.

SPREADER SETTINGS

To help obtain uniform distribution, apply one-half the required amount of product in one direction, then apply the balance at right angle.

Application Rate Lbs. per 1,000 Sq. Ft.	Lilly/ Miller	Central	Scotts PF-1	Ortho 84-65
2 1/2	4.75	7.0	5.5	1.75
3 1/2	5.75	7.5	6.5	3.0
5	7	9	7.5	3.5
10	12.5	11.5	10.5	6.5


TO HELP PREVENT INFESTATION OF BUILDINGS BY . . .

Ants, Boxelder Bugs, Crickets, Earwigs and Sowbugs (Pillbugs):
Treat a band of soil 6 to 10 feet around and adjacent to the building where pests are active and may find entrance. Use the same dosage as recommended for control of these pests on turf.

FOR ROOT MAINTENANCE CONTROL

IN DIRECT SEEDED CROPS: Apply only at seeding. Do not make more than one application per season.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Rutabagas*, Chinese Cabbage, Turnips*: Use 1/2 to 1 pound of product per 100 linear feet of row by placing over the seed in a 2 to 4 inch wide band at time of



planting. Sprinkle or irrigate with water after planting.

*With Rutabagas and Turnips, it would be realistic to expect only early season protection because of their extended growing season.

Restrictions: Do not apply more than 5 pounds of 1% CHLORBAN INSECT GRANULES per 1,000 sq. ft. Do not use rutabaga tops for food or feed purposes.

Radishes: Use 5 ounces of product per 100 linear feet of row by application in the seed furrow with the seed at planting time.

Restriction: Do not apply more than 6 pounds of 1% CHLORBAN INSECT GRANULES per 1,000 sq. ft.

1 cup of 1% CHLORBAN INSECT GRANULES weighs approximately 4 ounces.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals.

Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep out of reach of children. May be harmful if swallowed. Do not take internally. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air if symptoms of cholinesterase inhibition appear and get medical attention immediately.


If on Skin: Wash thoroughly with soap and water.

If in Eyes: Flush with plenty of water. Get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, by injection only, is antidotal.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.



Filled by weight not by volume. Product may settle after packaging.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DURSBAN is the Reg. T.M. of the Dow Chemical Co.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Reg. No. 802-560
EPA Est. 802-OR-1
30226

