102 171 FYX5

LILLY/MILLER
Casoron Granules

Controls Weeds & Grasses in Evergreens, Roses, Shrubs, Fruits & Nuts

ACTIVE INGREDIENT:

Dichlobenil: 2,6-Dichlorobenzonitrile . . . 1.5%

INERT INGREDIENTS 98.5%

Trademark and research discovery of Duphar, B.V., Amsterdam, Holland.

TREATS 1,600 SQ. FT.

Keep out of reach of children CAUTION
See back panel for additional cautions

NET WEIGHT 10 LBS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe dust. Do not allow contact with eyes or skin. In case of contact, flush eyes or skin with plenty of water. Get medical attention of irritation persists. Wash thoroughly after handling. Avoid contamination of feed and foodstuff.

ENVIRONMENTAL HAZARDS

Cover, collect, or incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

SAFETY RULES

IMPORTANT! Always wash hands, face, and arms with soap and water before, smoking, eating, drinking, or toileting. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is a danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP and get help right away.

CASORON GRANULES will selectively control annual grassy and certain perennial weeds around established ornamental trees and shrubs, fruit and nut trees, and berries, listed on this label. One application, made as directed, is generally enough to give season-long control.

CASORON GRANULES when applied to the soil as directed will give selective control of annual grassy and broadleaf weeds and certain perennial weeds such as:

Annual
Bluegrass
Artemesia
Bluegrass
Bull Thistle

Fiddleneck
Field Morning
Glory
Florida Purslane
(Pusley)

Purslane Quackgrass (from seed) Ragweed Red Deadnettle

ACCEPTED

JUN - 5 1939

Under the Federal Insecticide.
Fungicule, and Rodenticide Act
as disconded, for the pesticide
registered under
EPA Reg. No. 27

Camphorweed
Canaja Thistle
Carpetweed
Chickweed
Citron Melon
Coffeeweed
Crabgrass
Cudweed
Curly Dock
Dandelion
Dogfennel
Evening
Primrose
False
Dandelion

Foxtail
Gisekia
Groundsel
Henbit
Horsetail
Jerusalemoak
Goosefoot
Knotweed
Lambsquarters
Milkweed Vine
(Morrenia
odorata)
Natalgrass
Peppergrass
Pineapple Weed

Plantain

English Walnut

Redroot Pigweed
Rosarypea
Russian Knapweed
Russian Thistle
Shepherdspurse
Smartweed
Spanishneedles
Spurge
Teaweed
Texas Panicum
(Hurrahgrass)
Wild Carrot
Wild Mustard
Wild Radish
Yellow Woodsorrel

CASORON GRANULES are to be applied only to established woody plants - those which have been planted for at least four months.

CASORON GRANULES can be used for weed control on the following Ornamental Woody Shrubs:

Amur Corktree Arborvitae Ash Azalea 1 Barberry Beauty Bush Birch Box Elder Boxwood Camellia Caragana Cleyera Japonica Cotoneaster Cottonwood Crabapple Deutzia Dogwood Eastern Red Cedar Elm Elwood Cypress

Euonymous Filbert Flowering Almond Flowering Quince Forsythia Gardenia Goldenraintree Hackberry Heather Holly 2 Honeysuckle Ivy (Hedera Helix) Juniper Kinnikinnick Laurel Leucothoe Lilac Linden Locust Magnolia Maple

Mock Orange Mountain Ash Nandina Oak Osmanthus (Holly-Olive) Pachistima Pecan Photinia Pittosporum Poplar Privet Pyracantha Rhododendron Rockrose Rose Russian Olive: Squaw-carpet. Weigela Willow

1 Kurume, Mollis, Hardy Hybrid types and hardy native species ohly.
2 Except I. Crenata, I. Rotunda, I. Vomitoria.

CASORON GRANULES may be used for weed control on the following Edible Fruits & Nuts (except California):

Apple Blackberry Filbert Grape

Pear Prune

Yew (Taxus)

Blueberry Cherry Nectarine Peach Plum Raspberry

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before Application

1. Remove existing weeds.

2. Thoroughly work and smooth the area to be treated.

- 3. Carefilly measure area to be treated and determine the number of square
- 4. If area to be treated is large, it should be divided into smaller sections and each treated individually.

Application and Rate

1. Shake CASORON RANULES uniformly on the soil at the races specified.

FOR ANNUAL WEEDS: Apply in early spring before weeds emerge or after removing weeds.

FOR PERENNIAL WEEDS: For best control, apply after a killing frost from late fall to early winter.

2. ORNAMENTAL WOODY SHRUBS:

Use at rate of 1 ounce for each 10 sq. ft. to be treated. 10 pounds treats 1,600 sq. ft.

FRUIT TREES, NUTS, BLUBERRIES, GRAPES (Except California): Apply 1 - 1 1/2 ounce per 10 sq. ft. Use the higher rate - 1 1/2 ounce per 10 sq. ft. if weed infestation is heavy or if most weeds are perennial. Wait 30 days after making application before harvesting stone fruits or nuts.

BLACKBERRIES, RASPBERRIES (Except California): Use 1 ounce per 10 sq. ft. Use only on established plantings. Apply before new shoots emerge in spring.

3. Use the diagram below as a guide for uniform distribution of granules.

DIAGRAM

- 4. Rates higher than those indicated may cause injury.
- 5. More accurate and uniform application can be obtained by applying one-half of the required material over the entire treatment area and repeating the application with the remaining material...
- 6. CASORON GRANULES should not be used on sloping areas, by cause runoff may result in injury to lawn or other plants in adjacent areas.

After Application

- 1. Unless the soil is wet, sprinkle lightly to moisten the soil 1/4 to 1/2 inch deep. Do not flood, avoid puddling and washing.
- . Avoid disturbing soil in the treated area as long as weed control is

being maintained.

NOTE: Use only on the plants listed. Do not apply to transplant or cutting beds or to newly set transplants, and do not seed or transplant into treated areas for one year after application. Do not use on extremely light sandy soils. Do not plant rotational crops, on which Dichlobenil is not registered, in treated soil within one (1) year after application.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse bag (container). Securely wrap bag (container) in several layers of newspaper and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Filled by weight not by volume. Product may settle after packaging.

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218 EPA Reg. No. 802-571 EPA Est. 802-0R-1

