

Lilly/Miller
SUPER RICH FEED & WEED
With Trimec® Herbicide

TRIMEC is a very efficient weed killer. TRIMEC will kill many known broadleaf weeds infesting lawns*.

*Bahia, Bentgrass, Bermuda, Bluegrass, Fescue, Rye and Zoysia.

15-2-4

COVERAGE 2,500 SQ. FT.

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	0.28%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.13%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.03%
INERT INGREDIENTS	99.56%

*2,4-Dichlorophenoxyacetic acid equivalent 0.24%. Isomer Specific by AOAC Method

**2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 0.10%

***3,6-dichloro-o-anisic acid equivalent 0.02%

NET WT. 16.6 LBS.

BEST AVAILABLE COPY

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

3 WEED KILLERS GET MORE THAN JUST DANDELIONS

SUPER RICH FEED & WEED contains nutrients shown in this exploded view of this quality lawn care blend.

NITROGEN, water soluble

Just add water and it goes to work feeding your lawn immediately.

Nitrogen is a protein builder, helping grass to grow vigorous and strong. Gives your lawn its rich green color.

NITROGEN, water insoluble, organic

Lilly/Miller's special time-released nitrogen. Released to the grass roots slowly and continuously, by action of bacteria in the soil. Feeds your lawn over a long period of time. Builds protein and color through steady feeding.

PHOSPHORIC ACID

ACCEPTED

JUN 15 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 802-579

This important nutrient helps early growth and root formation. Helps young plants grow strong.

POTASH

Potash encourages a strong root system. Builds up your lawn's resistance to disease.

SULFUR

An essential element for protein movement. Helps nitrogen promote strong, sturdy plant growth.

IRON

Iron is necessary for chlorophyll production. This important element helps turn plants that deep, emerald green.

SUPER RICH FEED & WEED contains TRIMEC HERBICIDE, a combination of three proven weed killers. Together, they kill a wide range of broadleaf lawn weeds - dandelions, plantain, chickweed, knotweed, clover, Japanese clover, henbit, buttercup, spurge, lawn daisy, puncture vine, and most other common broadleaf weeds in Bahia, Bentgrass, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns.

SUPER RICH FEED & WEED not only feeds your lawn with five important plant nutrients, but kills lawn weeds - all in one step! It's the easy, time-saving way to get a vigorous, deep green, weed-free carpet of lush, thick grass.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Induce vomiting if swallowed. Wash hands after use. Keep out of reach of children. In case of contact flush with plenty of water. For eyes get medical attention. Avoid contact with skin, eyes and clothing. Wear long pants, long-sleeved shirt and shoes when applying this product. Avoid breathing dust. Do not reenter or permit others to reenter treated areas until granules have been washed into the soil and grass has dried.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEST AVAILABLE COPY

USE PRECAUTIONS: 2,4-D, MCPP and Dicamba are injurious to most plants other than grasses so this material should be applied carefully on a day with low wind velocity to avoid drift onto valuable plants. Do not use on St. Augustine grass, Dichondra, carpetgrass or clover lawns. May cause possible injury to bentgrass. Do not apply within the drip line of desirable trees and shrubs. Rinse spreader with clear water after use.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron for plant nutrition which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.


or

All products containing Iron - from Ferrous Sulfate, Fritted Micronutrients, etc. - have the potential to stain concrete, especially new or damp concrete in or near areas where the product can be tracked, thrown by broadcast spreader, or washed by water. Extra precautions must be taken to avoid such application or movement. Immediate removal of the product from concrete surfaces by blowing, sweeping and/or hosing will lessen the potential for staining.

Apply 16.6 lbs. to 2,500 sq. ft.
Apply any time weeds are growing actively-spring, summer or fall. Applications to moist foliage and soil, when temperature is between 50 degrees and 80 degrees F., are the most effective. Applications during hot weather (over 80 degrees F.) or on dry soil may cause some temporary damage to turf.

1. Do not apply to a new lawn. Wait until lawn is 3 months old.
2. Do not mow lawn for two days prior to application to have sufficient weed foliage to receive weed killers.
3. Apply to moist soil and foliage using an accurate spreader and correct dial setting from chart below. Make sure lawn is moist at time of application.
4. Do not water lawn for 24 hours after application. Resume watering, if needed, after 24 hours.
5. Make second application after one month if control of hard-to-kill weeds is not complete.

HOW TO SPREAD LAWN FERTILIZER



Lawns must be thoroughly watered the day before application. Begin spreading by laying down two strips of lawn food at each end of lawn. Fill in remaining area as shown here. Close the spreader when you stop walking or make turns to avoid uneven coverage. Stopping and starting with the spreader box open will waste lawn food. Overlap previous wheel track slightly, as shown, to avoid gaps in coverage. Wait 24 hours for herbicide to work. Then water in to start the greening process.

SPREADER SETTINGS

To calibrate your spreader, adjust spreader setting to apply 1 pound to 150 square feet of turf.

APPLICATION RATE	Lilly/Miller	Turf	Scotts Models	Spreader
		King	35 & 75	
16.6 lbs. per 2,500 sq. ft.	TO BE DETERMINED			

NOTICE

The spreader settings given on this bag have been carefully determined by actual test in new equipment. Incorrect dial settings, walking speed, wear, spreader calibration and general condition of spreader can materially affect the rate of flow. Since these conditions are beyond our control, neither The Chas. H. Lilly Co. nor the retail dealer can guarantee exact coverage as indicated by the recommended dial settings.

GUARANTEED FERTILIZER ANALYSIS:

- Total Nitrogen (N) 15%
 - 11.3% Ammoniacal Nitrogen
 - 0.25% Urea Nitrogen
 - 0.15% Other Water Soluble Nitrogen
 - 1.0% Coated Slow Release Urea Nitrogen
 - 2.3% Other Water Insoluble Nitrogen
- Available Phosphoric Acid (P2O5) 2%
- Soluble Potash (K2O) 4%

Primary Nutrients From:

Ammonium Phosphate; Ammonium Sulfate; Muriate of Potash; Processed Tankage; Sulfur-Coated Urea.

- Sulfur (S) 14%
- Iron (Fe) 2%

Secondary And Micronutrients From: Ammonium Sulfate; Iron Sulfate; Processed Tankage; Sulfur-Coated Urea.



or

GUARANTEED FERTILIZER ANALYSIS:

Total Nitrogen (N) 15%
 9.2% Ammoniacal Nitrogen
 0.9% Urea Nitrogen
 1.5% Slowly Available Water Soluble Nitrogen
 2.4% Coated Slow Release Urea Nitrogen
 1.0% Other Water Insoluble Nitrogen
 Available Phosphoric Acid (P2O5) 2%
 Soluble Potash (K2O) 4%

Primary Nutrients From:

Ammonium Phosphate; Ammonium Sulfate; Muriate of Potash; Methylene Ureas; Sulfur-Coated Urea; Gypsum.

Sulfur (S) 14%

Iron (Fe) 2%

Secondary And Micronutrients From: Ammonium Sulfate; Iron Sulfate; Processed Tankage; Sulfur-Coated Urea.

STORAGE AND DISPOSAL

Store this product in its original container out of reach of children and domestic animals. Do not reuse bag. Securely wrap original container in several layers of newspaper and discard in trash.

Filled by weight not by volume. Product may settle after packaging.

LIMIT OF WARRANTY AND LIABILITY

The Chas. H. LILLY Co. (LILLY) warrants that this product conforms to the chemical or other description on the label (and/or instructions) and is reasonably fit for the purposes set forth on the label (and/or instructions) when used strictly in accordance with the label (and/or instructions) under the conditions described there. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the following conditions:

*Buyer and all users shall promptly notify LILLY of any claims whether based in contract, negligence, strict liability, other tort, or otherwise.

*THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF LIABILITY OF LILLY FOR ANY LOSS, INJURY OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (including claims based upon contract, negligence, strict liability, other tort, or otherwise) SHALL BE:

1. The purchase price paid for the product, or at the election of LILLY
2. The replacement of such quantity of the product as is involved.

IN NO EVENT SHALL LILLY BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES.

BEST AVAILABLE COPY

*LILLY agents have no authority to give warranties or make representations on its behalf, concerning the use, quality or performance of this product, or in any other respect.

*The use, handling, storage, and other influencing factors are beyond the control of LILLY. Buyer assumes all risk of loss, injury, or damage due to use, handling, storage and other influencing factors, including but not limited to, incompatibility with other products, and weather conditions.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

TRIMEC is the reg. TM of PBI/Gordon Corporation.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Reg. No. 802-579
EPA Est. 802-WA-1

