

802-579

10/03/2002

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OCT 3 2002

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Molly Hickman
Central Garden & Pet DBA Lilly Miller Brands
P.O. Box 2289
Clackamas, OR 97015

Dear Ms. Hickman:

Subject: Revised Labels
Lilly/Miller Lawn Weed Killer
EPA Registration No. 802-485
Lilly/Miller Super Rich Feed & Weed ✓
EPA Registration No. 802-579
Lilly/Miller Ready-To-Use Lawn Weed Killer
EPA Registration No. 802-580
Your Submission Dated September 17, 2002

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- For each route of exposure in the First Aid section it is preferred that it be highlighted with bullets or in separate paragraphs. Refer to PR Notice 2001-1. A copy can be obtained at the following internet site:

http://www.epa.gov/opppmsd1/PR_Notices/pr2001-1.pdf

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records. If you have any questions concerning this letter, please contact Mr. James Stone at 703-305-7391.

Sincerely yours,

A handwritten signature in cursive script, appearing to read "Joanne".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

LILLY/MILLER
SUPER RICH FEED & WEED

A premium blend of natural and slow-release fertilizers for long-lasting green
With Trimec® broadleaf weed killer

15-5-10

Feeds 5,000 Sq. Ft.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

OCT 3 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
~~507-578~~

ACTIVE INGREDIENTS:

- *Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid 0.28%
- **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)
propionic acid 0.13%
- ***Dimethylamine salt of Dicamba
(3,6-dichloro-o-anisic acid) 0.03%
- INERT INGREDIENTS 99.56%

- *2,4-Dichlorophenoxyacetic acid equivalent 0.24%. Isomer Specific by AOAC Method
- **2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 0.10%
- ***3,6-dichloro-o-anisic acid equivalent 0.02%

Keep out of reach of children

CAUTION

See back panel for additional precautionary statements

NET WT. 33.3 LBS. (15.1 kg)

Your Lawn Can Be a Super Rich® Lawn in Just Four Easy Steps

Now there's a Super Rich Lawn Food for each season of the year. Start now by applying this season's recommended product. And then every three months, apply the appropriate Super Rich Lawn Food. It's as simple as that. By following the Super Rich annual program, your lawn will remain in peak condition all year long.

Winter: January – March

Super Rich Moss Control Lawn Food kills moss quickly so grass has room to grow. Greens and thickens your lawn for up to 3 months. For mild-winter climates where moss isn't a problem, apply Super Rich Fall & Winter Lawn Food.

Spring: April – June

Super Rich Feed & Weed with Trimec® controls dandelions, clover, plantain and many other broadleaf weeds while greening your lawn for up to 3 months.

Summer: July – September

Super Rich Lawn Food with 2% iron produces a dark green, deep-rooted lawn. Keeps on greening for up to 3 months. Now with non-staining iron to reduce rust spots on concrete and clothing.

Fall: October – December

Super Rich Fall & Winter Lawn Food with extra potassium keeps your lawn green through the fall and promotes thick, sturdy growth. Prepares your lawn for winter and helps it green up fast next spring.

Improved Super Rich lawn foods are now made with:

- 25% natural ingredients, including dried poultry waste and sulfate of potash magnesia (a naturally-occurring mineral)
- Four types of nitrogen for quick greening and long-term feeding
- University-endorsed ratios of nitrogen, phosphorus and potassium
- Non-staining iron* for a dark green lawn without the worry of rust spots on concrete and clothing

*In all Super Rich lawn foods except Super Rich® Moss Control Lawn Food.

Super Rich Feed & Weed kills weeds while greening and thickening your lawn. It kills dandelion, clover, black medic (Japanese clover), plantain, chickweed, buttercup, spurge, English daisy, and puncture vine. Super Rich Feed & Weed contains Trimec®—a combination of three systemic weed killers—to kill weeds from top to bottom.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep out of reach of children. Avoid contact with skin, eyes and clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Avoid breathing dust. Do not allow people (other than applicator) or pets on treatment area during application. Do not reenter or permit others to reenter treated areas until granules have been washed into the soil and grass has dried.

FIRST AID

Have product container or label with you when calling a poison control center or doctor, or going for treatment. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further

treatment advice. If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The maximum number of broadcast applications per treatment site is two per year. For household use only.

USE PRECAUTIONS

2,4-D, MCPP and Dicamba are injurious to most plants other than grasses, so this material should be applied carefully on a day with low wind velocity to avoid drift onto valuable plants. Do not use on St. Augustinegrass, dichondra, carpetgrass or clover lawns. May cause possible injury to bentgrass. Do not apply within the drip line of desirable trees and shrubs. Rinse spreader with clear water after use.

For best results, apply 33.3 lbs. to 5,000 sq. ft.

Apply any time weeds are growing actively-spring, summer or fall. Weed control is best when the foliage and soil are moist and the air temperature is between 50 and 80 degrees F. When applied during hot weather (over 80 degrees F) or on dry soil, turfgrass may be temporarily damaged.

1. Do not apply to a new lawn. Wait until lawn is 3 months old.
2. Do not mow the lawn for two days before application so that there are enough weed leaves to catch the weed killer granules.
3. Apply to moist soil and foliage using an accurate spreader. Make sure lawn is moist at time of application.
4. Do not water lawn for 24 hours after application. Resume watering, if needed, after 24 hours.
5. Make second application after one month if control of hard-to-kill weeds is not complete. Or use Lilly Miller® Lawn Weed Killer to control broadleaf weeds without harming your lawn.

SETTING YOUR LAWN SPREADER

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch the flow rate. If fertilizer appears to be coming out either slower or faster than you are covering the lawn, adjust the setting. Close the spreader when filling, turning or stopping.

Setting
(33.3 lbs. for 5,000 sq. ft.)

Drop Spreader Models

Earthway Ev-N-Spread	10
Red Devil/Precision	3
Scotts AccuGreen 1000	8.75
Scotts AccuGreen 2000	9
Scotts AccuGreen 3000	9.25
Scotts Precisiongreen (PF3)	8.5
Sta Green 4500	2.5
Vigoro 4000V	13

Rotary Spreader Models

These spreaders can toss lawn food a great distance. Do not allow granules to fall in flower or shrub beds or around trees. These plants can be hurt.

Earthway Ev-N-Spread	15
Red Devil/Precision 1000	9.5
Scotts SpeedyGreen 1000	9.5
Scotts SpeedyGreen 2000	10.5
Scotts SpeedyGreen 3000	9
Sta Green 4500	7
Vigoro SB 4000V	10

Captions for illustrations:

Lawns must be thoroughly watered the day before application. Then water again lightly, just prior to application. Begin spreading by laying down two strips of lawn food at each end of lawn. Fill in remaining area as shown here. Close the spreader when you stop walking or make turns to avoid uneven coverage. Stopping and starting with the spreader box open will waste lawn food. Overlap previous wheel track slightly, as shown, to avoid gaps in coverage. Wait 24 hours for herbicide to work. Then water lawn to start the greening process.

GUARANTEED FERTILIZER ANALYSIS:

- Total Nitrogen (N) 15%
- 1.35% Ammoniacal Nitrogen
- 1.05% Water Insoluble Nitrogen

- 12.6% Urea Nitrogen*
- Available Phosphate Acid (P2O5) 5%
- Soluble Potash (K2O) 10%
- Calcium (Ca) 2%
- Magnesium (Mg)2%
- Sulfur (S)6%
- Iron (Fe)2%
- *4.2% slowly available urea nitrogen from sulfur-coated urea.

Derived from dried poultry waste, urea, sulfur-coated urea, ammonium sulfate, monoammonium phosphate, sulfate of potash magnesia, muriate of potash, iron sucrate.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.wa.gov/agr/>.

PROTECT YOUR PETS

This fertilizer contains animal manures and/or byproducts that may attract dogs or other pets. Consumption of this product may make animals sick. To minimize potential problems: 1) Store away from pets, 2) Thoroughly water in as directed.

STORAGE AND DISPOSAL

Store this product in its original container and keep out of reach of children and domestic animals. If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Filled by weight not by volume. Product may settle after packaging.

TRIMEC is the reg. TM of PBI/Gordon Corporation.

SATISFACTION GUARANTEED: If for any reason you are not satisfied with this product, send proof of purchase to the address shown and we will gladly refund your purchase price.

LILLY MILLER BRANDS
 PO BOX 2289
 CLACKAMAS, OREGON 97015
 EPA Reg. No. 802-579
 EPA Est. 35497-OR-1
 www.lillymiller.com
 EPA draft 9/17/2002

