





UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Rebecca M. Horton Central Garden & Pet dba Lilly Miller Brands 1340 Treat Blvd., Suite 650 Creek, CA 94597

AUG 13 2009

Dear Ms. Horton:

Subject: Label Amendment According to Agency Letter (dated May, 29, June 9 and 24, 2009)

Liller Miller Super Rich Feed & Weed

EPA Reg. No. 802-579

Your Submission Dated July 6, 2009

The label, referred to above, submitted in connection with the application under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the following change:

- 1. Move the emergency phone number statement to the bottom of the First Aid section.
- 2. Remove the word "over" so the sentence reads "To prevent product run-off, do not water the treated area to the point of runoff or apply when raining or when rain is expected that day." This is located at the bottom of page 2 and top of page 3, the last sentence in the last paragraph of the Environmental Hazards section.
- 3. Be sure to express the following per MCPP-p RED, the following restrictions must be added to the directions for use:

"For broadcast treatments:

Limited to 2 applications per year. Maximum of 0.75 lb ae per acre per application. Minimum of 30 days between applications. page 2 EPA Reg. No. 802-579

A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(3). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions, please contact Tracy White by phone at (703) 308-0042 or via email at white.tracy@epa.gov.

Sincerely yours,

Joanne I. Miller Product manager (23) Herbicide Branch Registration Division (7505P)

Enclosure

LILLY/MILLER SUPER RICH FEED & WEED x-x-x

[Additional Brand Name: Lilly Miller Super Rich Feed & Weed Lawn Food 15-5-10]

Optional Marketing Statements:

A premium blend of natural and slow-release fertilizers for long-lasting green With Trimec® broadleaf weed killer

Feeds 5,000 Sq. Ft.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	0.28%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	0.06%
Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.03%
OTHER INGREDIENTS:	99.63%
Total	00.00%

This product contains:

0.24% 2,4-Dichlorophenoxyacetic acid equivalent 0.05% (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 0.02% 3,6-dichloro-o-anisic acid equivalent Isomer Specific by AOAC Methods.

ACCEPTED with COMMENTS In EPA Letter Dated:

AUG 13 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No

Keep out of reach of children

802-579

CAUTION

See back panel for Additional Precautionary Statements and First Aid.

NET WT. 33.3 LBS. (15.1 kg)

EPA' Ēši. 35497-OR-1 Page 1 of 6 pages

EPA Reg. No. 802-579 July 6, 2009 - amend

FIRST AID

Have product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact [1-800-265-0761] [the National Pesticide Information Center at 1-800-858-7378] for emergency medical treatment information.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. When using this product, wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estimates, bays and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter or street back onto the treated area. To prevent

EPA Est. 35497-OR-1 Page 2 of 6 pages product run-off do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household use only.

General Precautions and Restrictions

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not enter or allow others (including children and pets) to enter the treated areas (except those involved with the watering) until the watering-in is completed and the surface is dry. The maximum number of applications per treatment site is two per year. Do not exceed recommended application rate. Allow 30 days between applications.

2,4-D, MCPP and Dicamba are injurious to most plants other than grasses. Do not use on St. Augustinegrass, dichondra, carpetgrass or clover lawns. May cause possible injury to bentgrass. Do not apply within the drip line of desirable trees and shrubs. Do not apply to a new lawn; wait until lawn is 3 months old. Rinse spreader with clear water after use.

Apply any time weeds are growing actively-spring, summer or fall. Weed control is best when the foliage and soil are moist and the air temperature is between 50 and 80 degrees F. When applied during hot weather (over 80 degrees F) or on dry soil, turfgrass may be temporarily damaged.

- 1. Apply 33.3 lbs. to 5,000 sq. ft.
- 2. Do not mow the lawn for two days before application so that there are enough weed leaves to catch the weed killer granules.
- 3. Apply to moist soil and foliage using an accurate spreader. Make sure lawn is moist at time of application.
- 4. Do not water lawn for 24 hours after application. Resume watering, if needed, after 24 hours.
- 5. Make second application after one month if control of hard-to-kill weeds is not complete. Or use Lilly Miller® Lawn Weed Killer to control broadleaf weeds without harming your lawn.

EPA Est: 3:

EPA Est. 35497-OR-1 Page 3 of 6 pages

EPA Reg. No. 802-579 July 6, 2009 - amend

SETTING YOUR LAWN SPREADER

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch the flow rate. If fertilizer appears to be coming out either slower or faster than you are covering the lawn, adjust the setting. Close the spreader when filling, turning or stopping.

Setting
(33.3 lbs. for 5,000 sq. ft.)
10
3
8.75
9
9.25
8.5
2.5
13

Rotary Spreader Models

These spreaders can toss lawn food a great distance. Do not allow granules to fall in flower or shrub beds or around trees. These plants can be hurt.

Earthway Ev-N-Spread	15
Red Devil/Precision 1000	9.5
Scotts SpeedyGreen 1000	9.5
Scotts SpeedyGreen 2000	10.5
Scotts SpeedyGreen 3000	9
Sta Green 4500	7
Vigoro SB 4000V	10

Captions for illustrations:

Lawns must be thoroughly watered the day before application. Then water again lightly, just prior to application. Begin spreading by laying down two strips of lawn food at each end of lawn. Fill in remaining area as shown here. Close the spreader when you stop walking or make turns to avoid uneven coverage. Stopping and starting with the spreader box open will waste lawn food. Overlap previous wheel track slightly, as shown, to avoid gaps in coverage. Wait 24 hours for herbicide to work. Then water lawn to start the greening process.

GUARANTEED ANALYSIS:

Total Nitrogen (N) 15% 2.9% Ammoniacal Nitrogen 11.1% Urea Nitrogen* 1.0% Water Insoluble Nitrogen

> EPA Est. 35497-OR-1 Page 4 of 6 pages

Available Phosphate (P2O5) 5% Soluble Potash (K2O) 10% Calcium (Ca) 2% Magnesium (Mg)2% Sulfur (S)6% Iron (Fe)2%

Derived from ammonium sulfate, urea, polymer coated urea, dried poultry waste, ammonium phosphate, sulfate of potash magnesia, muriate of potash, iron sucrate.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm.

PROTECT YOUR PETS

This fertilizer contains animal manures and/or byproducts that may attract dogs or other pets. Consumption of this product may make animals sick. To minimize potential problems: 1) Store away from pets, 2) Thoroughly water in as directed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

<u>PESTICIDE STORAGE</u>: Store this product in its original container and keep out of reach of children and domestic animals. <u>PESTICIDE DISPOSAL</u>: If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. <u>CONTAINER DISPOSAL</u>: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

Optional Marketing Statements

Your Lawn Can Be a Super Rich ® Lawn in Just Four Easy Steps

Now there's a Super Rich Lawn Food for each season of the year. Start now by applying this season's recommended product. And then every three months, apply the appropriate Super Rich Lawn Food. It's as simple as that. By following the Super Rich annual program, your lawn will remain in peak condition all year long.

Winter: January - March

Super Rich Moss Control Lawn Food kills moss quickly so grass has room to grow. Greens and thickens your lawn for up to 3 months. For mild-winter climates where moss isn't a problem, apply Super Rich Fall & Winter Lawn Food.

Spring: April – June

Super Rich Feed & Weed with Trimec® controls dandelions, clover, plantain and many other broadleaf weeds while greening your lawn for up to 3 months.

Summer: July – September

EPA Est. 35497-OR-1 Page 5 of 6 pages

))) 00

^{*4.2%} slowly available urea nitrogen from polymer coated urea.

Super Rich Lawn Food with 2% iron produces a dark green, deep-rooted lawn. Keeps on greening for up to 3 months. Now with non-staining iron to reduce rust spots on concrete and clothing.

Fall: October - December

Super Rich Fall & Winter Lawn Food with extra potassium keeps your lawn green through the fall and promotes thick, sturdy growth. Prepares your lawn for winter and helps it green up fast next spring.

Improved Super Rich lawn foods are now made with:

- 25% natural ingredients, including dried poultry waste and sulfate of potash magnesia (a naturally-occurring mineral)
- Four types of nitrogen for quick greening and long-term feeding
- University-endorsed ratios of nitrogen, phosphorus and potassium
- Non-staining iron* for a dark green lawn without the worry of rust spots on concrete and clothing

*In all Super Rich lawn foods except Super Rich® Moss Control Lawn Food.

Super Rich Feed & Weed kills weeds while greening and thickening your lawn. It kills dandelion, clover, black medic (Japanese clover), plantain, chickweed, buttercup, spurge, English daisy, and puncture vine. Super Rich Feed & Weed contains Trimec®--a combination of three systemic weed killers—to kill weeds from top to bottom.

Filled by weight not by volume. Product may settle after packaging.

TRIMEC is the reg. TM of PBI/Gordon Corporation.

SATISFACTION GUARANTEED: If for any reason you are not satisfied with this product, send proof of purchase to the address shown and we will gladly refund your purchase price.

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

[Lot no. to be added at production]

LILLY MILLER BRANDS 1340 Treat Blvd., Suite 650 Walnut Creek, CA 94597 www.lillymiller.com

Lilly Miller is a registered trademark of Central Garden & Pet.

