

LILLY/MILLER
HOSE 'N GO WEED & FEED 15-0-0

Contains TRIMEC

Control weeds and feed grass in one application. Just attach to a garden hose and spray.

Ready to use. Covers up to 3,750 sq. ft.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	5.673%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	2.738%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.629%
INERT INGREDIENTS	90.96%

- +*0.44 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 4.708%
- **0.21 lb. 2-(2-methyl-4-chlorophenoxy)propionic acid per gallon or 2.272%
- ***0.048 lb. Dicamba (3,6-dichloro-o-anisic acid) per gallon or 0.522%

+Isomer specific by AOAC method.
TRIMEC is the Reg. T.M. of PBI/Gordon Corporation.

Keep out of reach of children
CAUTION
See back panel for additional precautionary statements.

NET CONTENTS 24 FL. OZ.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Avoid contamination of food or feedstuffs.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water when disposing of equipment washwaters. Do not contaminate domestic or irrigation waters.

STOP. READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. On new lawns, wait until the grass has been mowed two or three times and has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow 1 to 2 days before treating. Do not water lawn within 24 hours after treatment.

WHERE TO USE: HOSE 'N GO WEED & FEED is especially effective on all hard-to-kill weeds such as Poison Ivy, Poison Oak, Sheep Sorrel and Spurge. It also controls most other broadleaf weeds, including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

HOW TO USE: Shake well before using. 1. Connect spray nozzle to garden hose. Be sure that valve is in "off" position. 2. Remove bottle cap and screw bottle onto nozzle. Turn on water. 3. Switch valve to "on" position to start flow of water. Use valve to shut off flow of water at any time. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to determine remaining area to be covered).

OTHER USE PRECAUTIONS: Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85 degrees F. Seed can be safely sown 3 weeks after application. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, closely-mowed bentgrass, or on lawns or turf where desirable clovers are present. Do not apply to surfaces above the root zone of trees and ornamentals. Do not exceed specified dosages for any area, do not apply within the dripline of trees and ornamentals.

HOSE 'N GO WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) 15%
15% Urea Nitrogen
Primary nutrients from Urea.

STORAGE AND DISPOSAL